




PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : C07D 213/85, A61K 31/44		A1	(11) International Publication Number: WO 00/47561
			(43) International Publication Date: 17 August 2000 (17.08.00)
(21) International Application Number: PCT/SE99/01403		Marmorsvägen, S-752 44 Uppsala (SE). WALLBERG, Hans [SE/SE]; Mangårdsvägen 10, S-141 51 Huddinge (SE). SAHLBERG, Christer [SE/SE]; Mälarhöjdsvägen 5L, S-129 40 Hägersten (SE).	
(22) International Filing Date: 18 August 1999 (18.08.99)			
(30) Priority Data: 09/249,317 12 February 1999 (12.02.99) US PCT/SE99/00194 15 February 1999 (15.02.99) SE			
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/249,317 (CIP) Filed on 12 February 1999 (12.02.99)			
(71) Applicant (for all designated States except US): MEDIVIR AB [SE/SE]; Lunastigen 7, S-141 44 Huddinge (SE).			
(72) Inventors; and (75) Inventors/Applicants (for US only): ZHOU, Xiao-Xiong [SE/SE]; Kallkärsvägen 12, S-141 41 Huddinge (SE). JOHANSSON, Nils-Gunnar [SE/SE]; Bäverstigen 19, S-150 23 Enhörna (SE). WÄHLING, Horst [DE/SE]; Tempelriddarevägen 3, S-127 61 Skärholmen (SE). SUND, Christian [FI/SE]; Bäckgårdsvägen, S-143 21 Vårby (SE). SALVADOR, Lourdes [PH/SE]; Saltängsgatan 1B, S-602 22 Nörrebro (SE). LINDSTRÖM, Stefan [SE/SE];		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
		Published With international search report.	

(54) Title: NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS



## (57) Abstract

Non-nucleoside reverse transcriptase inhibitors of formula (P-1) wherein: Ar1 is an unsaturated, optionally substituted, mono or bicyclic ring structure comprising 0 to 3 hetero atoms selected from S, O and N; Ar2 is an aromatic, optionally substituted, monocyclic ring structure comprising at least one nitrogen hetero atom and zero to two further hetero atoms selected from S, O and N; R4 and R5 are independently H or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl, C<sub>1</sub>-C<sub>5</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkanoyloxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, amino, carboxy, carbamoyl, cyano, halo, hydroxy, aminomethyl, hydroxymethyl, carboxymethyl, or halo substituted C<sub>1</sub>-C<sub>6</sub> alkyl mercapto, nitro; or R4 and R5 join to form a 3-6 membered, optionally substituted ring structure; R6 is O or S; Rx is the residue of a natural or unnatural amino acid; and L\* is a linker moiety which is ether-, carbonate- or ester-bound to the adjacent oxygen and ester linked to Rx; and pharmaceutically acceptable salts thereof are anti-HIV agents with favourable pharmacokinetic properties.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						


Non-nucleoside reverse transcriptase inhibitorsTechnical field

This invention relates to non-nucleoside reverse transcriptase inhibitors active  
5 against HIV-1 and having improved pharmacokinetic properties. The invention  
further relates to the synthesis of such compounds and their use in antiviral  
methods and compositions.

Background to the invention

10 Non nucleoside reverse transcriptase inhibitors (NNRTI) bind to an allosteric  
site on reverse transcriptase and represent an important development in the  
arsenal of drugs against HIV, particularly HIV-1. International patent  
application WO 93/03022, discloses thiourea NNRTI, later denoted PETT  
(phenyl ethyl thiazolyl thiourea) compounds in J Med Chem 39 6 1329-1335  
15 (1995) and J Med Chem 39 21 4261-4274 (1996). International patent  
application no WO95/06034 discloses urea isosteres of PETT NNRTIs.  
PCT/SE99/00053 published after the priority and international filing dates of  
the present application discloses particularly active urea NNTRI compounds  
with the formula:

20



wherein

R8 is halo;

R9 is C<sub>1</sub>-C<sub>3</sub> alkyl;

25 R10 is halo, especially bromo or cyano.

The contents of each of the above documents are specifically incorporated by  
reference.


Although the urea and thiourea NNRTI disclosed in the above documents are extremely active against reverse transcriptase, especially that of HIV-1, the nature of the HIV virus with its extreme lack of replicative fidelity and consequent tendency to rapid resistance development prompts a demand for further antiretroviral agents. Additionally, modern HIV therapy regimes, denoted HAART, Highly Active Anti Retroviral Therapy, administer antivirals as combinations of three or more antivirals of various classes, which combinations are administered for prolonged periods, if not for life. HAART requires the patient to follow a complicated dosing schedule with sometimes dozens of tablets per day taken at various times of the day in some cases before and in other cases after the ingestion of food. There is thus a need for antiretroviral preparations allowing greater flexibility in dosing to facilitate patient compliance.

To pursuit of such flexibility of dosing, the abovementioned PCT/SE99/00053 proposes certain prodrugs of the compounds of formula P2 above building on the phenolic hydroxy group. These prodrugs comprise aminobenzoyl or pyridyl esters of this phenolic hydroxy, wherein the amino group may be optionally substituted with an amino acid peptidically bound thereto.

#### Brief description of the invention

We have now discovered that a particular class of prodrug provides enhanced pharmacokinetic performance in respect of NNRTIs of the PETT class.

Accordingly a first aspect of the invention provides compounds of the formula P1:



P-1

wherein:

Ar1 is an unsaturated, optionally substituted, mono or bicyclic ring structure comprising 0 to 3 hetero atoms selected from S, O and N;

Ar2 is an aromatic, optionally substituted, monocyclic ring structure comprising at least one nitrogen hetero atom and zero to two further hetero

5 atoms selected from S, O and N;

R4 and R5 are independently H, a substituent selected from C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl, C<sub>1</sub>-C<sub>5</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkanoyloxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, amino, carboxy, carbamoyl, cyano, halo, hydroxy, aminomethyl, hydroxymethyl, carboxymethyl, or halo substituted C<sub>1</sub>-C<sub>6</sub> alkyl mercapto, nitro,  
10 or R4 and R5 join to form a 3 – 6 membered , optionally substituted ring structure;

R6 is O or S;

L\* is a linker moiety which is ether-, carbonate- or ester-bound to the adjacent oxygen and ester linked to Rx;

15 Rx is the residue of a natural or unnatural amino acid;  
and pharmaceutically acceptable salts thereof.

Favoured Ar1 groups include phenyl, pyridyl, naphthyl, indanyl, indolyl, quinolyl, isoquinolyl, benzopyridyl, benzoimidazol, benzothiazolyl, benzopyrazinyl,  
20 benzotriazolyl, benzopyrimidyl, benzopyridazinyl, purinyl, pyrazinyl, pyrrolyl pyrazolyl, pyrimidinyl, pyridazinyl, triazolyl, cyclohexenyl and the like, especially pyrid-2-yl, pyrid-3-yl and phenyl. The –O-L\*-Rx moiety is conveniently located at the 2 or 5 position of ring Ar1 relative to the linkage to the rest of the molecule.

25

Favoured Ar2 groups include pyridyl, pyrimidinyl, pyrrolyl, pyrazinyl, pyridazinyl, triazolyl, thiazolyl, thiadiazolyl and the like, especially pyridyl. Preferably Ar2 comprises a nitrogen atom at the 2-position relative to the linkage to the (thio)urea. The optional substituent to Ar2, if present, is  
30 preferably located para to the linkage to the (thio)urea.

R6 is preferably oxygen, thus defining a urea PETT derivative.

R4 and R5 as a ring structure can comprise optionally substituted cyclopropyl, cyclobutyl, cyclopentyl or the like, preferably cis-cyclopropyl. Alternatively R4 and R5 are conveniently H.

- 5 The optional substituents described herein conveniently comprise up to three substituents selected from C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl, C<sub>2</sub>-C<sub>8</sub> alkenoxy, C<sub>1</sub>-C<sub>6</sub> alkoxy C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkanoyl, haloC<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkanoyloxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, amino, carboxy, carbamoyl, cyano, halo, hydroxy, aminomethyl, carboxymethyl, hydroxymethyl, nitro, aryl,
- 10 substituted (as herein defined) aryl, or -SO<sub>2</sub>D or -C(=O)D where D is C<sub>1</sub>-C<sub>6</sub> alkyl, aryl, substituted (as herein defined) aryl or amino;

The optional substituent to Ar2 preferably comprises halo, especially bromo or cyano. The optional substituent(s) to Ar1 comprise one or two members

15 independently selected from the group consisting H, fluoro, bromo, chloro, hydroxy, methoxy, ethoxy, amino, dimethylamino, acetyl, propionyl, butyryl and the like. Where Ar2 is a six membered ring, the optional substituents are conveniently located at positions 2 and/or 6 relative to the linkage to the rest of the molecule.


20

- C<sub>1</sub>-C<sub>n</sub> alkyl includes such groups as methyl, ethyl, n-propyl, isopropyl, n-butyl, s-butyl, t-butyl, n-pentyl, n-hexyl, 3-methyl pentyl and the like. The term halo refers to chloro, bromo, fluoro and iodo. C<sub>1</sub>-C<sub>n</sub> alkoxy refers to groups such as methoxy, ethoxy, propoxy, t-butoxy and the like. C<sub>2</sub>-C<sub>n</sub> alkenyl, refers to
- 25 groups such as vinyl, 1-propen-2-yl, 1-buten-4-yl, 1-penten-5-yl, 1-buten-1-yl and the like. C<sub>1</sub>-C<sub>n</sub> alkylthio includes methylthio, ethylthio, t-butylthio and the like. C<sub>1</sub>-C<sub>n</sub> alkanoyloxy includes acetoxo, propionoxo, formyloxy, butyryloxy and the like. C<sub>2</sub>-C<sub>n</sub> alkenoxy includes ethenyloxy, propenyloxy, isobutoxyethenyl and the like. HaloC<sub>1</sub>-C<sub>n</sub> alkyl includes alkyls as defined herein
- 30 substituted 1 to 3 times by a halogen including trifluoromethyl, 2-dichloroethyl, 3,3-difluoropropyl and the like. The term amine includes groups such as NH<sub>2</sub>, NHMe, N(Me)<sub>2</sub> which may optionally be substituted with halogen, C<sub>1</sub>-C<sub>7</sub> acyloxy, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, nitro, carboxy, carbamoyl, carbamoyloxy, cyano, methylsulphonylamino and the like. Carboxy, carboxymethyl and

carbamoyl include the corresponding pharmaceutically acceptable C<sub>1</sub>-C<sub>6</sub> alkyl and aryl esters.

Particularly preferred compounds of the present invention have the formula

5 P-2:



wherein

R8 is halo, especially halo;

10 R9 is C<sub>1</sub>-C<sub>3</sub> alkyl;

R10 is halo or cyano;


and L\*Rx is as defined above.

A preferred subset of compounds within Formula P2, particularly with regard  
15 to pharmacokinetics, has R10 as cyano. A further favoured subset of compounds within Formula P2, particularly with regard to ease of forming prodrugs, comprise compounds wherein R10 is bromo.

Preferably R8 is chloro and more preferably fluoro. Suitable R9 groups include methyl, isopropyl, n-propyl and preferably ethyl.

20

As depicted in Formula P2, the cyclopropyl ring is in the *cis* configuration, allowing two enantiomers, 1S, 2S and 1R, 2R



25 Each of these enantiomers are potent antiretrovirals, although the different enantiomers can display subtle differences in physiological properties. For


instance the 1S, 2S and 1R,2R enantiomers can show a different pattern of metabolism within the P450 system. The 1S,2S enantiomer of compounds wherein Rp10 is cyano is particularly preferred as it appears unique in being able to avoid key components of the P450 system. Other retroviral agents  
5 such as the HIV protease inhibitor ritonavir interact extensively with the P450 system, leading to an array of undesirable physiological responses including extensive alteration of the metabolism of other co-administered drugs. This is of particular concern with pharmaceuticals administered for a chronic infection where patients can expect to take a number of pharmaceuticals for years, if  
10 not decades.

Preferred NNRTI mother compounds for preparing prodrugs in accordance with the invention thus include:

- (1S, 2S)-N-[*cis*-2-(6-fluoro, 2-hydroxy, 3-propionylphenyl)-cyclopropyl]-N'-(5-  
15 cyanopyrid-2-yl)-urea,  
(1S, 2S)-N-[*cis*-2-(6-fluoro, 2-hydroxy, 3-butyrylphenyl)-cyclopropyl]-N'-(5-  
cyanopyrid-2-yl)-urea,  
(1S, 2S)-N-[*cis*-2-(6-fluoro, 2-hydroxy, 3-acetylphenyl)-cyclopropyl]-N'-(5-  
cyanopyrid-2-yl)-urea,  
20 (1S, 2S)-N-[*cis*-2-(6-fluoro, 2-hydroxy, 3-propionylphenyl)-cyclopropyl]-N'-(5-  
bromopyrid-2-yl)-urea,  
(1S, 2S)-N-[*cis*-2-(6-fluoro, 2-hydroxy, 3-butyrylphenyl)-cyclopropyl]-N'-(5-  
bromopyrid-2-yl)-urea,  
(1S, 2S)-N-[*cis*-2-(6-fluoro, 2-hydroxy, 3-acetylphenyl)-cyclopropyl]-N'-(5-  
25 bromopyrid-2-yl)-urea,  
and the corresponding 1R, 2R enantiomers.

As the compounds of formula P1 and in particular P2 generally include an electron withdrawing group on the phenol/heteroaryl ring to which the  
30 prodrug moiety is attached, it is generally preferred to avoid a simple ester linkage for L\*. Accordingly L\* preferably comprises an ether or carbonate linkage to the phenolic (or other heteroaryl) hydroxy of the mother compound. A particularly convenient ether linkage has the formula:






Where PEST is the dehydroxy residue of the NNRTI, the arrowed oxygen is the phenolic (or other heterocyclic) hydroxy, Ry and Ry' are independently H or C<sub>1</sub>-C<sub>3</sub> alkyl and A is the esterified amino acid residue Rx. Alternatively, A can comprise an intermediate linking moiety to which Rx is esterified. Examples of such an intermediate linking moiety include a straight or branched chain hydroxycarboxylic acid derivative, such as 3-hydroxypropionate, 4-hydroxybutyrate and in particular L-lactic acid, wherein the carboxy function is esterified to the non-arrowed oxygen above and the hydroxy function is esterified to the amino acid. The intermediate linking moiety may comprise an aryl or heteroaryl component bearing an (alkylenyl)carboxy function amenable for esterification with the amino acid of Rx. The form and preparation of linking groups suitable for use in the compounds of the invention is extensively described in our copending application no PCT/SE99/00194, the contents of which are incorporated by reference.

Alternatively A as an intermediate linking moiety can comprise a carbonate linkage to the alkyloxy group depicted above, in which case A further comprises a branched or straight chain hydroxycarboxylic acid derivative amenable to esterification with the amino acid of Rx. Similarly a carbonate linkage bearing A moiety may comprise an aryl or heteroaryl component bearing an (alkylenyl)carboxy function amenable for esterification with the amino acid of Rx.

Thus a convenient group of prodrugs within the scope of the invention include those of the formula P3:



wherein


- 5 R8, R9, R10 are as described above;  
 Ry and Ry' are independently H or C<sub>1</sub>-C<sub>3</sub> alkyl;  
 Rx is the ester residue of a natural or unnatural amino acid;  
 and pharmaceutically acceptable salts thereof.
- 10 Typically both of Ry and Ry' are H. Rx is preferably the ester residue of an aliphatic amino acid such as alanine, tertiary leucine or leucine, preferably isoleucine and especially valine, Preferably Rx is derived from an L-amino acid.
- 15 Preferred compounds within Formula P3 thus include;  
 (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(L-valyloxymethyloxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea,  
 (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(L-isoleucyloxymethyloxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea,  
 20 (1R, 2R)-N-{ *cis*-2-[6-fluoro-2-(L-valyloxymethyloxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea,  
 (1R, 2R)-N-{ *cis*-2-[6-fluoro-2-(L-isoleucyloxymethyloxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea,  
 (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(L-valyloxymethyloxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-bromopyridyl)]urea,  
 25 (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(L-isoleucyloxymethyloxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-bromopyridyl)]urea,

(1R, 2R)-N-{*cis*-2-[6-fluoro-2-(L-valyloxymethoxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-bromopyridyl)]urea,

(1R, 2R)-N-{*cis*-2-[6-fluoro-2-(L-isoleucyloxymethoxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-bromopyridyl)]urea,

5 and pharmaceutically acceptable salts thereof.

An alternative preferred group of prodrugs of the invention have the Formula P4:




P4

where R8, R9, R10, Ry and Ry' are as defined above. L and R<sub>2</sub> define a linker group and residue of an amino acid, such as those depicted in copending application PCT/SE99/00194.

Typically both of Ry and Ry' are H. Rx is preferably the ester residue of an aliphatic amino acid such as alanine, tertiary leucine or leucine, preferably isoleucine and especially valine. Preferably Rx is derived from an L-amino acid.

Favoured compounds within formula P4 include those of the formula P5:




P5

where R<sup>8</sup> R<sup>9</sup>, R<sup>10</sup>, R<sub>y</sub>, R<sub>y</sub>' and R<sub>x</sub> are as defined above and Alk<sup>b</sup> is C<sub>1</sub>-C<sub>6</sub> optionally branched, optionally monounsaturated alkyl.

Favoured compounds within Formula P5 thus include:

- 5 (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(2,2-dimethyl-3-(L-valyloxy)-propionyl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,
- (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(2-methyl-3-(L-valyloxy)-propionyl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,
- (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(3-(L-valyloxy)-propionyl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,
- 10 (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(2,2-dimethyl-3-(L-valyloxy)-propionyl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-bromopyridyl)] urea,
- (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(2-methyl-3-(L-valyloxy)-propionyl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-bromopyridyl)] urea,
- 15 (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(3-(L-valyloxy)-propionyl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-bromopyridyl)] urea,
- (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(2,2-dimethyl-3-(L-isoleucyloxy)-propionyl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,
- (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(2-methyl-3-(L-isoleucyloxy)-propionyl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,
- 20 (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(3-(L-isoleucyloxy)-propionyl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,
- (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(2,2-dimethyl-3-(L-isoleucyloxy)-propionyl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-bromopyridyl)] urea,
- 25 (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(2-methyl-3-(L-isoleucyloxy)-propionyl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-bromopyridyl)] urea,
- (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(3-(L-isoleucyloxy)-propionyl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-bromopyridyl)] urea,
- (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(4-(L-valyloxy)-butyryl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,
- 30 (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(4-(L-isoleucyloxy)-butyryl-oxymethyloxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,

- (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(4-(L-valyloxy)-butyryl-oxymethoxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-bromopyridyl)] urea,  
 (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(4-(L-isoleucyloxy)-butyryl-oxymethoxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-bromopyridyl)] urea,  
 5 (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2-(L-valyloxy)-propionyl-oxymethoxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,  
 (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2-(L-isoleucyloxy)-propionyl-oxymethoxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,  
 (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2-(L-valyloxy)-propionyl-oxymethoxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-bromopyridyl)] urea,  
 10 (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2-(L-isoleucyloxy)-propionyl-oxymethoxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-bromopyridyl)] urea,  
 and the corresponding (1R,2R) enantiomers thereof.
- 15 One variant of a branched Alk<sup>b</sup> in Formula P5 can be substituted with hydroxy which in turn is esterified with a further Rx, thus defining a compound of the Formula P6:




P6

- 20 where R8, R9, R10, Alk, Ry, Ry', and Rx are as defined above, RZ is H or C<sub>1</sub>-C<sub>3</sub> alkyl ( ) represents a methylene group and m and n are independently 0, 1 or 2.

- Particularly favoured values for Alk, m and n include: methylene:1:1 and  
 25 absent: 1:0 respectively.

A further favoured group of compounds has the Formula P7:



P7


where R8, R9, R10, Alk, Ry, Ry', m, n Rz and Rx are as defined above or

5 wherein the  $(\text{---})_m\text{---O---R}_2$  arm is absent.

Typically both of Ry and Ry' are H. Preferably Rz is H. Rx is preferably the  
 ester residue of an aliphatic amino acid such as alanine, tertiary leucine or  
 leucine, preferably isoleucine and especially valine, Preferably Rx is derived  
 10 from an L-amino acid.

Particularly favoured values for Alk, m and n include:absent:1:1, thus defining  
 a glycerol derivative.

15 Where the  $(\text{---})_m\text{---O---R}_2$  arm is absent in formula P7, the compounds have the  
 structure P7':




P7'

Convenient values for Alk and n include absent:1 with Ry, Ry and Rz' as H.

Favoured compounds within Formula P-7 thus include

- (1S,2S)-N-[*cis*-2-(6-fluoro-2-(1,3-bis-L-valyloxy-2-(oxycarbonylmethoxy)propyl)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,
- 5 (1S,2S)-N-[*cis*-2-(6-fluoro-2-(1,3-bis-L-isoleucyloxy-2-(oxycarbonylmethoxy)propyl)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,
- (1S,2S)-N-[*cis*-2-(6-fluoro-2-(1,3-bis-L-valyloxy-2-(oxycarbonylmethoxy)propyl)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,
- 10 (1S,2S)-N-[*cis*-2-(6-fluoro-2-(1,3-bis-L-isoleucyloxy-2-(oxycarbonylmethoxy)propyl)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,
- (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(2-(L-valyloxy)-ethoxycarbonyloxymethyloxy)-3-propionylphenyl]]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,
- 15 (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(2-(L-isoleucyloxy)-ethoxycarbonyloxymethyloxy)-3-propionylphenyl]]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,
- (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(2-(L-valyloxy)-ethoxycarbonyloxymethyloxy)-3-propionylphenyl]]cyclopropyl}-N'-[2-(5-bromopyridyl)] urea,
- 20 (1S, 2S)-N-{ *cis*-2-[6-fluoro-2-(2-(L-isoleucyloxy)-ethoxycarbonyloxymethyloxy)-3-propionylphenyl]]cyclopropyl}-N'-[2-(5-bromocyanopyridyl)] urea,
- and the corresponding R,R enantiomers.

- 25 A further favoured group of compounds omit the alkyleneoxy group immediately adjacent the phenolic (or other heteroaryl) hydroxy function of the compound of formula P1 or P2. An example of such compounds has the formula P8:



P8

where R8, R9, R10, Rx, and Alk<sup>b</sup> are as defined above.

Currently favoured values for Alk<sup>b</sup> include methylene, ethylene, 1,1-dimethylethylene, 2,2-isopropylene, butylene and, in the case of said -ORx substitution described below, glycerol.


Favoured compounds within formula P-8 thus include:

- (1S,2S)-N-[*cis*-2-(6-fluoro-2-(L-valyloxymethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,
- (1S,2S)-N-[*cis*-2-(6-fluoro-2-(L-isoleucyloxymethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,
- (1S,2S)-N-[*cis*-2-(6-fluoro-2-(L-valyloxymethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,
- (1S,2S)-N-[*cis*-2-(6-fluoro-2-(L-isoleucyloxymethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,
- (1S,2S)-N-[*cis*-2-(6-fluoro-2-(2-(L-valyloxy)ethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,
- (1S,2S)-N-[*cis*-2-(6-fluoro-2-(2-(L-isoleucyloxy)ethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,
- (1S,2S)-N-[*cis*-2-(6-fluoro-2-(2-(L-valyloxy)ethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,
- (1S,2S)-N-[*cis*-2-(6-fluoro-2-(2-(L-isoleucyloxy)ethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,
- (1S,2S)-N-[*cis*-2-(6-fluoro-2-(3-(L-valyloxy)propoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,
- (1S,2S)-N-[*cis*-2-(6-fluoro-2-(3-(L-isoleucyloxy)propoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,



- (1*S*,2*S*)-*N*-[*cis*-2-(6-fluoro-2-(3-(*L*-valyloxy)propoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-*N'*-[2-(5-bromopyridyl)]urea,  
 (1*S*,2*S*)-*N*-[*cis*-2-(6-fluoro-2-(3-(*L*-isoleucyloxy)propoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-*N'*-[2-(5-bromopyridyl)]urea,  
 5 (1*S*,2*S*)-*N*-[*cis*-2-(6-fluoro-2-(4-(*L*-valyloxy)butoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-*N'*-[2-(5-cyanopyridyl)]urea,  
 (1*S*,2*S*)-*N*-[*cis*-2-(6-fluoro-2-(4-(*L*-isoleucyloxy)butoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-*N'*-[2-(5-cyanopyridyl)]urea,  
 (1*S*,2*S*)-*N*-[*cis*-2-(6-fluoro-2-(4-(*L*-valyloxy)butoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-*N'*-[2-(5-bromopyridyl)]urea,  
 10 (1*S*,2*S*)-*N*-[*cis*-2-(6-fluoro-2-(4-(*L*-isoleucyloxy)butoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-*N'*-[2-(5-bromopyridyl)]urea,  
 and the corresponding *R*, *R* enantiomers

- 15 As with Formula P5/P6 and P7/P7', Alk<sup>b</sup> in formula P8 can comprise an additional -O-R<sub>z</sub> substitution to define a compound of the formula P8'



P8'

where each of the variables is as defined above.

20

- The invention further provides pharmaceutical compositions comprising the compounds of formula P1 and pharmaceutically acceptable carriers or diluents therefor. Additional aspects of the invention provide methods for the inhibition of HIV comprising administering a compound of the formula I to a  
 25 subject afflicted with HIV. The invention also extends to the use of the compounds of formula I in therapy, such as in the preparation of a medicament for the treatment of HIV infections.

In treating conditions caused by HIV, the compounds of formula I are preferably administered in an amount to achieve a plasma level of the compounds of Formula P1 of around 10 to 1000 nM and more preferably 100 to 500 nM. This corresponds to a dosage rate, depending on the  
 5 bioavailability of the formulation, of the order 0.01 to 10 mg/kg/day, preferably 0.1 to 2 mg/kg/day. A typical dosage rate for a normal adult will be around 0.05 to 5 g per day, preferably 0.1 to 2 g such as 500-750 mg, in one to four dosage units per day.


10 In keeping with the usual practice with HIV inhibitors it is advantageous to co-administer one to three additional antivirals to provide synergistic responses and to ensure complementary resistance patterns. Such additional antivirals may include AZT, ddI, ddC, D4T, 3TC, abacavir, adefovir, adefovir dipivoxil, bis-POC-PMPA, foscarnet, hydroxyurea, Hoechst-Bayer HBY 097,  
 15 efavirenz, trovirdine, nevirapine, delaviridine, PFA, H2G (tamociclovir), ABT 606 (valtamociclovir stearate) DMP-450, loviride, ritonavir, saquinavir, indinavir, amprenavir (Vertex VX 478), nelfinavir and the like, typically at molar ratios reflecting their respective activities and bioavailabilities. Generally such ratio will be of the order of 25:1 to 1:25, relative to the compound of formula I.

20

Compounds of the invention are typically prepared by alkylation of the corresponding mother compounds of Formula P1 or especially P2, which are prepared by conventional means, such as the methodology described in WO95/06034 or PCT/SE99/00053.

25


In particular, the preparation of compounds of formula P-3 or P-4 generally proceeds by alkylation using conventional coupling conditions of a compound of the formula P-2 with the corresponding intermediate:



30

where  $R_y$ ,  $R_y'$  and  $L$  are as defined above and  $R_x$ -PG is  $R_x$  as defined, but N-protected with a conventional N-protecting group. Preferably the halogen activating group is iodo, which is in turn prepared by iodination of the corresponding chloro analogue. Typical coupling conditions include treatment with a base in an organic solvent such as THF prior to addition of the activated (halogenated) intermediate followed by conventional deprotection of the  $R_x$  N-protecting group.

Compounds of formula P8 are generally prepared by esterification of a compound of the formula P2 with an intermediate of the formula:



where  $\text{Alk}^{\text{b}*}$  is a functionalised  $\text{Alk}^{\text{b}}$  as described above, for example chloromethyl chloroformate, in an organic solvent, followed by iodination of the terminal chloro with NaI (or other activation of the functionalising group) and esterification/alkylation with an N-protected  $R_x$ .

After coupling of the activated  $L^*$ -(N-protected)- $R_x$  group to the mother NNRTI (or the two step coupling of the activated  $L^*$  moiety followed by coupling of the N-protected  $R_x$ ) the N-protecting group on the amino acid  $R_x$  is removed by conventional techniques such as palladium catalysed reduction or treatment with an organic acid such as trifluoroacetic acid.

Typical intermediates for a one step coupling thus comprise  
 iodomethyloxy-N-CBz-valyl,  
 iodomethyloxy-N-Boc-valyl,  
 iodomethyloxy-N-Fmoc-valyl  
 iodomethyloxy-N-CBz-isoleucyl,  
 iodomethyloxy-N-Boc-isoleucyl,  
 iodomethyloxy-N-Fmoc-isoleucyl,  
 iodomethyloxy-N-CBz-tertiaryleucyl,

iodomethyloxy-N-Boc-tertiaryleucyl,  
iodomethyloxy-N-Fmoc-tertiaryleucyl,  
and corresponding derivatives bearing other N-protecting groups.

- 5 Further typical intermediates for a one step coupling include  
2,2-dimethyl-3-(N-Boc-L-valyloxy)propionic acid iodomethyl ester  
3,3- bis (N-CBz-L-valyloxymethyl)-propionic acid iodomethyl ester,  
2-(N-CBz-L-valyloxy)ethoxycarbonyloxymethyl iodide  
iodomethyl 1,3-bis(N-benzyloxycarbonyl-L-valyloxy)-2-propyl carbonate,  
10 iodomethyl 2-methyl-2-(N-benzyloxycarbonyl-L-valyloxymethyl) propionate,  
iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxy)-DL-propionate.  
iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxy)isobutyrate.  
iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxy)-3-methyl-(S)-(+)-butyrate.  
iodomethyl 2-O-(N-benzyloxycarbonyl-L-valyloxy)-2-phenyl-DL-acetate  
15 iodomethyl 4-(N-benzyloxycarbonyl-L-valyloxy) benzoate.  
iodomethyl 5-(N-CBz-L-valyloxy)-2,2-dimethylvalerate  
2-(N-CBz-L-valyloxy)-ethyl iodomethyl carbonate  
4-(N-CBz-L-valyloxy) butyric acid iodomethyl ester  
iodomethyl-3-(N-benzyloxycarbonyl-L-valyloxy)-benzoate  
20 iodomethyl-3-(N-benzyloxycarbonyl-L-valyloxy)-propionate  
1,3-bis(N-tert-butoxycarbonyl-L-valyloxy)-2-propyl 1-iodoethyl carbonate  
3-(N-benzyloxycarbonyl-L-valyloxy)-2,2-dimethylpropyl iodomethyl carbonate  
iodomethyl 3,4-di-(N-CBZ-L-valyloxy)hydrocinnamate  
3-(N-CBZ-L-valyloxy)phenyl iodomethyl carbonate  
25 iodomethyl 2-(N-CBZ-L-valyloxy)phenylacetate  
iodomethyl 4-(N-CBZ-L-valyloxy)phenylacetate  
iodomethyl 4-(2-N-benzyloxycarbonyl-L-valyloxyethyl) benzoate  
iodomethyl 4-(N-benzyloxycarbonyl-L-valyloxy)cyclohexanoate.  
iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-ethyl butyrate  
30 2-(N-(iodomethoxycarbonyl)-amino)-2-methyl-1-(N-benzyloxycarbonyl-  
L-valyloxy)-propane  
1-(2-N-CBz-L-valyloxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid  
iodomethyl ester  
iodomethyl 5-[(N-benzyloxycarbonyl-L-valyloxy)methyl]-2-furoate

- Iodomethyl 4-(2-N-benzyloxycarbonyl-L-valyloxyethoxy)-benzoic acid  
2,2-dimethyl-3-(N-Boc-L-isoleucyloxy)propionic acid iodomethyl ester  
3,3- bis (N-CBz-L-isoleucyloxymethyl)-propionic acid iodomethyl ester,  
2-(N-CBz-L-isoleucyloxy)ethoxycarbonyloxymethyl iodide
- 5 Iodomethyl 1,3-bis(N-benzyloxycarbonyl-L-isoleucyloxy)-2-propyl carbonate,  
Iodomethyl 2-methyl-2-(N-benzyloxycarbonyl-L-isoleucyloxymethyl)  
propionate,  
Iodomethyl 2-(N-benzyloxycarbonyl-L-isoleucyloxy)-DL-propionate.  
Iodomethyl 2-(N-benzyloxycarbonyl-L-isoleucyloxy)isobutyrate.
- 10 Iodomethyl 2-(N-benzyloxycarbonyl-L-isoleucyloxy)-3-methyl-(S)-(+)-butyrate.  
Iodomethyl 2-(N-benzyloxycarbonyl-L-isoleucyloxy)-2-phenyl-DL-acetate  
Iodomethyl 4-(N-benzyloxycarbonyl-L-isoleucyloxy) benzoate.  
Iodomethyl 5-(N-CBz-L-isoleucyloxy)-2,2-dimethylvalerate  
2-(N-CBz-L-isoleucyloxy)-ethyl iodomethyl carbonate
- 15 4-(N-CBz-L-isoleucyloxy) butyric acid iodomethyl ester  
Iodomethyl-3-(N-benzyloxycarbonyl-L-isoleucyloxy)-benzoate  
Iodomethyl-3-(N-benzyloxycarbonyl-L-isoleucyloxy)-propionate  
1,3-bis(N-tert-butoxycarbonyl-L-isoleucyloxy)-2-propyl 1-iodoethyl carbonate  
3-(N-benzyloxycarbonyl-L-isoleucyloxy)-2,2-dimethylpropyl iodomethyl  
20 carbonate  
Iodomethyl 3,4-di-(N-CBz-L-isoleucyloxy)hydrocinnamate  
3-(N-CBz-L-isoleucyloxy)phenyl iodomethyl carbonate  
Iodomethyl 2-(N-CBz-L-isoleucyloxy)phenylacetate  
Iodomethyl 4-(N-CBz-L-isoleucyloxy)phenylacetate
- 25 Iodomethyl 4-(2-N-benzyloxycarbonyl-L-isoleucyloxyethyl) benzoate  
Iodomethyl 4-(N-benzyloxycarbonyl-L-isoleucyloxy)cyclohexanoate,  
Iodomethyl 2-(N-benzyloxycarbonyl-L-isoleucyloxymethyl)-2-ethyl butyrate,  
2-(N-(iodomethoxycarbonyl)-amino)-2-methyl-1-(N-benzyloxycarbonyl-  
L-isoleucyloxy)-propane,
- 30 1-(2-N-CBz-L-isoleucyloxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid  
iodomethyl ester  
Iodomethyl 5-[(N-benzyloxycarbonyl-L-isoleucyloxy)methyl]-2-furoate  
Iodomethyl 4-(2-N-benzyloxycarbonyl-L-isoleucyloxyethoxy)-benzoic acid  
and the corresponding chloromethyl analogues.

Although the above list describes Boc and CBz N-protecting groups on the amino acid, it will be appreciated that other N-protecting groups are also available.

- 5 Copending international application no PCT/SE99/00194 comprehensively describes the preparation of analogous intermediates.

The term "N-protecting group" or "N-protected" as used herein refers to those groups intended to protect the N-terminus of an amino acid or peptide or to  
10 protect an amino group against undesirable reactions during synthetic procedures. Commonly used N-protecting groups are disclosed in Greene, "Protective Groups in Organic Synthesis" (John Wiley & Sons, New York, 1981), which is hereby incorporated by reference. N-protecting groups include acyl groups such as formyl, acetyl, propionyl, pivaloyl, t-butylacetyl, 2-  
15 chloroacetyl, 2-bromoacetyl, trifluoroacetyl, trichloroacetyl, phthalyl, o-nitrophenoxyacetyl,  $\alpha$ -chlorobutyryl, benzoyl, 4-chlorobenzoyl, 4-bromobenzoyl, 4-nitrobenzoyl, and the like; sulfonyl groups such as benzenesulfonyl, p-toluenesulfonyl, and the like, carbamate forming groups such as benzyloxycarbonyl, p-chlorobenzyloxycarbonyl,  
20 p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, 2-nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl, 3,4-dimethoxybenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitro-4,5-dimethoxybenzyloxycarbonyl, 3,4,5-trimethoxybenzyloxycarbonyl, 1-(p-biphenyl)-1-methylethoxycarbonyl,  $\alpha,\alpha$ -dimethyl-3,5-  
25 dimethoxybenzyloxycarbonyl, benzhydryloxycarbonyl, t-butoxycarbonyl, diisopropylmethoxycarbonyl, isopropylloxycarbonyl, ethoxycarbonyl, methoxycarbonyl, allyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, phenoxycarbonyl, 4-nitrophenoxycarbonyl, fluorenyl-9-methoxycarbonyl, cyclopentylloxycarbonyl, adamantylloxycarbonyl, cyclohexylloxycarbonyl,  
30 phenylthiocarbonyl, and the like; alkyl groups such as benzyl, triphenylmethyl, benzyloxymethyl and the like; and silyl groups such as trimethylsilyl and the like. Favoured N-protecting groups include formyl, acetyl, benzoyl, pivaloyl,

t-butylacetyl, phenylsulfonyl, benzyl, t-butoxycarbonyl (BOC) and benzyloxycarbonyl (Cbz).

Hydroxy and/or carboxy protecting groups are also extensively reviewed in  
5 Greene *ibid* and include ethers such as methyl, substituted methyl ethers such as methoxymethyl, methylthiomethyl, benzyloxymethyl, t-butoxymethyl, 2-methoxyethoxymethyl and the like, silyl ethers such as trimethylsilyl (TMS), t-butyltrimethylsilyl (TBDMS) tribenzylsilyl, triphenylsilyl, t-butyltriphenylsilyl triisopropyl silyl and the like, substituted ethyl ethers such as 1-ethoxymethyl,  
10 1-methyl-1-methoxyethyl, t-butyl, allyl, benzyl, p-methoxybenzyl, diphenylmethyl, triphenylmethyl and the like, aralkyl groups such as trityl, and pixyl (9-hydroxy-9-phenylxanthene derivatives, especially the chloride). Ester hydroxy protecting groups include esters such as formate, benzylformate, chloroacetate, methoxyacetate, phenoxyacetate, pivaloate,  
15 adamantate, mesitoate, benzoate and the like. Carbonate hydroxy protecting groups include methyl vinyl, allyl, cinnamyl, benzyl and the like.

The compounds of the invention can form salts which form an additional aspect of the invention. Appropriate pharmaceutically acceptable salts of the  
20 compounds of Formula I include salts of organic acids, especially carboxylic acids, including but not limited to acetate, trifluoroacetate, lactate, gluconate, citrate, tartrate, maleate, malate, pantothenate, isethionate, adipate, alginate, aspartate, benzoate, butyrate, digluconate, cyclopentanate, glucoheptanate, glycerophosphate, oxalate, heptanoate, hexanoate, fumarate, nicotinate,  
25 palmoate, pectinate, 3-phenylpropionate, picrate, pivalate, proprionate, tartrate, lactobionate, pivalate, camphorate, undecanoate and succinate, organic sulphonic acids such as methanesulphonate, ethanesulphonate, 2-hydroxyethane sulphonate, camphorsulphonate, 2-naphthalenesulphonate, benzenesulphonate, p-chlorobenzenesulphonate and p-toluenesulphonate;  
30 and inorganic acids such as hydrochloride, hydrobromide, hydroiodide, sulphate, bisulphate, hemisulphate, thiocyanate, persulphate, phosphoric and sulphonic acids. The compounds of the invention I may in some cases be isolated as the hydrate.

While it is possible for the active agent to be administered alone, it is preferable to present it as part of a pharmaceutical formulation. Such a formulation will comprise the above defined active agent together with one or more acceptable carriers or excipients and optionally other therapeutic ingredients. The carrier(s) must be acceptable in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient.

The formulations include those suitable for rectal, nasal, topical (including buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration, but preferably the formulation is an orally administered formulation. The formulations may conveniently be presented in unit dosage form, e.g. tablets and sustained release capsules, and may be prepared by any methods well known in the art of pharmacy.

Such methods include the step of bringing into association the above defined active agent with the carrier. In general, the formulations are prepared by uniformly and intimately bringing into association the active agent with liquid carriers or finely divided solid carriers or both, and then if necessary shaping the product. The invention extends to methods for preparing a pharmaceutical composition comprising bringing a compound of Formula I or its pharmaceutically acceptable salt in conjunction or association with a pharmaceutically acceptable carrier or vehicle. If the manufacture of pharmaceutical formulations involves intimate mixing of pharmaceutical excipients and the active ingredient in salt form, then it is often preferred to use excipients which are non-basic in nature, i.e. either acidic or neutral. Formulations for oral administration in the present invention may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active agent; as a powder or granules; as a solution or a suspension of the active agent in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water in oil liquid emulsion and as a bolus etc.




With regard to compositions for oral administration (e.g. tablets and capsules), the term suitable carrier includes vehicles such as common excipients e.g. binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, polyvinylpyrrolidone (Povidone), methylcellulose, ethylcellulose, sodium carboxymethylcellulose, hydroxypropylmethylcellulose, sucrose and starch; fillers and carriers, for example corn starch, gelatin, lactose, sucrose, microcrystalline cellulose, kaolin, mannitol, dicalcium phosphate, sodium chloride and alginic acid; and lubricants such as magnesium stearate, sodium stearate and other metallic stearates, stearic acid, glycerol stearate, silicone fluid, talc waxes, oils and colloidal silica. Flavouring agents such as peppermint, oil of wintergreen, cherry flavouring or the like can also be used. It may be desirable to add a colouring agent to make the dosage form readily identifiable. Tablets may also be coated by methods well known in the art. A tablet may be made by compression or moulding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active agent in a free flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface-active or dispersing agent. Moulded tablets may be made by moulding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may be optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active agent.

Other formulations suitable for oral administration include lozenges comprising the active agent in a flavoured base, usually sucrose and acacia or tragacanth; pastilles comprising the active agent in an inert base such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active agent in a suitable liquid carrier.

### Detailed Description

#### Preparation of intermediates

#### Example AA-I-1

2,3-Bis-(N-CBz-L-valyloxy)-propionic acid.

## a) t-Butyl 2,3-bis (N-CBz-L-valyloxy)propionate.

To a solution of t-butyl 2,3-dihydroxypropionate (2.43g, 15 mmole), N-CBz-L-valine (7.54g, 30 mmole) and DMAP (0.37g, 3 mmole) in 150 ml dichloromethane was added DCC (7.2g 35 mmole) and the mixture was stirred for two days at room temperature. The mixture was cooled to about 5°C and the urethane was filtered. The filtrate was evaporated, ethyl acetate was added and the organic phase washed twice with 5% acetic acid, 5% sodium hydrogen carbonate and water. The organic phase was dried with sodium sulfate filtered and evaporated under reduced pressure. The product was isolated by silica gel column chromatography. Yield: 8.2g = 86%


## b) 2,3-Bis-(N-CBz-L-valyloxy)-propionic acid.

To a solution of t-butyl 2,3-bis-(N-CBz-L-valyloxy)-propionate (7.2g, 11.4 mmole) in dichloromethane (25 ml) was added trifluoroacetic acid (25 ml) and the solution was stirred for five hours at room temperature. The solution was evaporated under reduced pressure and coevaporated two times with toluene. The product was isolated by silica gel column chromatography. Yield : 5.9g = 90% The intermediate may be esterified to the phenolic hydroxy group of the NNRTI using conventional esterification (ie a coupling agent such as DCC/DMAP but is preferably used as an intermediate linking moiety in conjunction with an methyloxy moiety as described above.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) 0.92 (m, 12H) 2.08 (m, 2H) 3.92-4.17 (m, 2H) 4.30-4.67 (m, 2H) 5.04 (s, 4H) 5.28 (m, 1H) 7.32 (m, 10H) 7.70 (m, 2H)

## Example AA-I-2

(S)-(+)-2-(N-CBz-L-valyloxy)propionic acid



a) 4-Methoxybenzyl (S) (+)-2-hydroxypropionate.

To a stirred solution of (S)(+)-2 hydroxypropionic acid (9.0g, 100 mmole) in  
 5 100 ml dry DMF was added potassium tert-butoxide (12.34g, 110 mmole) and  
 the mixture was stirred for one hour at 25°C. 4-Methoxybenzyl chloride (18.8g  
 120 mmole) was added and the mixture was stirred for six hours at 60°C. The  
 mixture was evaporated under reduced pressure and 250 ml ethyl acetate  
 was added. The organic phase was washed four times with water. The  
 10 organic phase was dried with sodium sulfate and concentrated in vacuo.  
 Yield: 15.6g = 74%

b) 4-Methoxybenzyl (S)-(+)-2-(N-CBz-L-valyloxy)propionate.

To a solution of 4-methoxybenzyl (S)-(+)-2-hydroxypropionate (7.6g, 36  
 mmole), N-CBZ-L-valine (10.05g, 40 mmole) and DMAP (0.98g, 8 mmole) in  
 15 150 ml dichloromethane was added a solution of DCC (8.3g, 40 mmole) and  
 the mixture was stirred overnight at room temperature. The mixture was  
 cooled to about 5°C and the urethane was filtered. The filtrate was evaporated  
 and the product was isolated by silica gel column chromatography. Yield:  
 14.4g = 90%


20 c) (S)-(+)-2-(N-CBz-L-valyloxy)propionic acid.

To a solution of 4-methoxybenzyl (S)-(+)-2-(N-CBz-L-valyloxy)propionate  
 (14.0g, 31.5 mmole) in dichloromethane (50 ml) was added trifluoroacetic acid  
 (25 ml) and the solution was stirred for five hours at room temperature. The  
 solution was evaporated under reduced pressure and coevaporated two times  
 25 with toluene. The product was isolated by silica gel column chromatography.  
 Yield: 9.4g = 92%. The intermediate may be esterified to the phenolic hydroxy  
 group of the NNRTI using conventional esterification (ie a coupling agent such  
 as DCC/DMAP but is preferably used as an intermediate linking moiety in  
 conjunction with an methyloxy moiety as described above.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) 0.94 (m, 6H) 1.46 (d, 3H) 2.12 (m, 1H) 4.05 (m, 1H) 4.92 (m, 1H) 5.06 (s, 2H) 7.34 (m, 5H) 7.68 (d, 1H)

Example AA-I-3

5 3,3-Bis (N-CBz-L-valyloxymethyl)-propionic acid



a) 4,4-bis (N-CBZ-L-valyloxymethyl)-but-1-ene.

10 To a solution of 2-allyl-1,3-propanediol (2.32g, 20 mmole), N-CBZ-L-valine (10.06g, 40 mmole) and DMAP (0.488g, 4 mmole) in 120ml dichloromethane was added DCC (9.08g, 44 mmole) in portions and the mixture was stirred overnight at room temperature. The mixture was cooled to 5°C and the urethane was filtered. The filtrate was evaporated and the product was  
15 isolated by silica gel column chromatography. Yield : 9.0g

b) 3,3-Bis (N-CBZ-L-valyloxymethyl)-propionic acid.

To a cooled solution of 4,4-bis (N-CBZ-L-valyloxymethyl)-but-1-ene (14.6g, 25 mmole) and tetrabutylammonium bromide (1.3g, 4 mmole) in 120ml benzene was added 100ml water. Under strong stirring potassium permanganate  
20 (15.8g, 100 mmole) was added in portions and the mixture was stirred for 2 hours between 15°C and 20°C . A sodium bisulfite aqueous solution was added to the slurry until the mixture was discolored. The mixture was acidified with 2N hydrochloric acid and extracted four times with ethyl acetate. The organic phase was washed two times with water, dried with sodium sulfate  
25 and evaporated under reduced pressure . The product was isolated by silica gel column chromatography. Yield: 7.5g . The intermediate may be esterified to the phenolic hydroxy group of the NNRTI using conventional esterification (ie a coupling agent such as DCC/DMAP but is preferably used as an intermediate linking moiety in conjunction with an methoxy moiety as  
30 described above.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.89 (m, 12H) 2.05 (m, 2H) 2.46 (m, 2H) 2.62 (m, 1H) 4.20 (m, 6H) 5.11 (s, 4H) 5.30 (m, 2H) 7.35 (m, 10H)

5 Example AA-I-4.

2-(N-CBZ-L-valyloxy)-propionic acid

a) 4-methoxybenzyl 2-hydroxypropionate.

To a stirred solution of DL -2 hydroxypropionic acid (9.0g, 100 mmole) in 100  
10 ml dry DMF was added potassium tert-butoxide (12.34g, 110 mmole) and the  
mixture was stirred for one hour at 60°C. 4-methoxybenzyl chloride (18.8g  
120 mmole) was added and the mixture was stirred for eight hours at 60°C.  
The mixture was evaporated under reduced pressure and 250 ml ethyl  
acetate was added. The organic phase was washed four times with water.  
15 The organic phase was dried with sodium sulfate and concentrated in vacuo.  
Yield: 16.8g

b) 4-methoxybenzyl 2-(N-CBZ-L-valyloxy)propionate.

To a solution of 4-methoxybenzyl 2-hydroxypropionate (4.2g, 20 mmole), N-  
CBZ-L-valine (5.02g, 20 mmole) and DMAP (0.24g, 2 mmole) in 100 ml  
20 dichloromethane was added a solution of DCC (4.54g, 22 mmole) and the  
mixture was stirred overnight at room temperature. The mixture was cooled to  
5°C and the urethane was filtered. The filtrate was evaporated and the  
product was isolated by silica gel column chromatography. Yield: 7.9g


c) 2-(N-CBZ-L-valyloxy)-propionic acid.

25 To a solution of 4-methoxybenzyl 2-(N-CBZ-L-valyloxy)-propionate (7.8g, 17.5  
mmole) in dichloromethane (100 ml) was added trifluoroacetic acid (10 ml)  
and the solution was stirred for one hour at room temperature. The solution  
was evaporated under reduced pressure and the product was isolated by  
silica gel column chromatography. Yield: 5.0g. The intermediate may be  
30 esterified to the phenolic hydroxy group of the NNRTI using conventional  
esterification (ie a coupling agent such as DCC/DMAP but is preferably used  
as an intermediate linking moiety in conjunction with a methoxy moiety as  
described above.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.94 (m, 6H) 1.56 (d, 3H) 2.30 (m, 1H) 4.42 (m, 1H) 5.12-5.30 (m, 4H) 7.28 (m, 5H)

5 Example AA-I-5

Succinic acid 2,3-bis-(N-CBZ-L-valyloxy)propyl ester



a) 4-Methoxybenzyl succinate monoester.

10 To a mixture of succinic anhydride (75g, 750 mmole) and 4-methoxybenzyl alcohol (69.1g, 500 mmole) in 1,4-dioxane (300ml) was added pyridine (79.1g, 1000 mmole) and the mixture was stirred for five hours at 80°C. The mixture was evaporated under reduced pressure and 600 ml of ethyl acetate and 60 ml of acetic acid were added. The organic phase was washed three times with  
15 water, dried with sodium sulfate and evaporated under reduced pressure. The product was recrystallized from toluene. Yield: 104 g.

b) Succinic acid 2,3-dihydroxy-propyl ester, 4-methoxybenzyl ester.

To a solution of glycerol (23.0g, 250 mmole), 4-methoxybenzyl succinate monoester (5.96 g, 25 mmole) and DMAP (0.36g, 3 mmole) in DMF (200ml)  
20 was added DCC (6.2g 30 mmole) and the mixture was stirred overnight at room temperature. The mixture was evaporated under reduced pressure and 150ml dichloromethane was added. The mixture was filtered and the solution washed twice with water. The water phase was extracted two times with dichloromethane and the combined organic phases were dried with sodium  
25 sulfate. The solution was evaporated under reduced pressure and the product was isolated by silica gel column chromatography.

Yield: 3.0g

c) Succinic acid 2,3-bis-(N-CBZ-L-valyloxy)-propyl ester, 4-methoxybenzyl ester.

To a stirred solution of succinic acid 2,3-dihydroxy-propyl ester, 4-methoxybenzyl ester (2.9g, 9.28 mmole), N-CBZ-L-valine (5.03g, 20 mmole) and DMAP (0.244g, 2 mmole) in dichloromethane (60ml) was added DCC (4.5g, 22 mmole) and the mixture was stirred overnight at room temperature.

- 5 The mixture was filtered and the solution was evaporated under reduced pressure. The product was isolated by silica gel column chromatography. Yield: 2.5g


d) Succinic acid 2,3-*bis*-(N-CBZ-L-valyloxy)propyl ester.

- To a solution of the above intermediate (2.3g, 2.95 mmole) in  
 10 dichloromethane (25ml) was added trifluoroacetic acid (2.5ml) and the solution was stirred for two hours at roomtemperature. The solution was evaporated under reduced pressure and the product was isolated by silica gel column chromatography. Yield: 1.8g .  
 The intermediate may be esterified to the phenolic hydroxy group of the  
 15 NNRTI using conventional esterification (ie a coupling agent such as DCC/DMAP but is preferably used as an intermediate linking moiety in conjunction with an methyloxy moiety as described above.

- <sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.92 (m, 12H) 2.12 (m, 2H) 2.64 (m, 4H) 4.32 (m, 4H) 5.10  
 20 (s, 4H) 5.22-5.50 (m, 3H) 7.34 (m, 10H)

#### Example AA-I-6

##### Succinic acid 1,3-*bis*-(N-CBZ-L-valyloxy)-2-propyl ester



25

- a) Succinic acid 1,3-dibromo-2-propyl ester, 4-methoxybenzyl ester.  
 To a solution of 1,3-dibromopropan-2-ol (21.8g, 100 mmole), succinic acid 4-methoxybenzyl ester (28.6g, 120 mmole) and DMAP (1.22g, 10 mmole) in  
 30 dichloromethane (400ml) was added DCC (24.8g, 120 mmole) in portions at about 10°C. The mixture was stirred overnight at room temperature and

cooled to about 5°C. The mixture was filtered and the solution was evaporated under reduced pressure. 600ml of ethyl acetate was added and the organic phase was washed twice with 5% acetic acid, 5% sodium hydrogen carbonate and water.

- 5 The solution was dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography. Yield: 34.8g.

b) Succinic acid 1,3-*bis*-(N-CBZ-L-valyloxy)-2-propyl ester, 4-methoxybenzyl ester.

- 10 To a solution of N-CBZ-L-valine ( 58.5 g, 232.8 mmole) in dried DMF (300ml) was added potassium-*tert*.-butoxide (24,68 g, 220 mmole) and the mixture was stirred for one hour at room temperature. A solution of succinic acid 1,3-dibromo-2-propyl ester, 4-methoxybenzyl ester (34 g, 77.6 mmole) in dried DMF (50ml) was added and the mixture was stirred for eighteen hours at  
15 60°C. The potassium bromide was filtered and the solution was evaporated under reduced pressure. 600ml of ethyl acetate was added and the organic phase washed twice with 5% sodium hydrogen carbonate and with water. The organic phase was dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column  
20 chromatography. Yield: 45g

c) Succinic acid 1,3-*bis*-(N-CBZ-L-valyloxy)-2-propyl ester.

- To a cooled solution of the intermediate immediately above (44.5 g, 57.1 mmole) in dichloromethane (500ml) was added trifluoroacetic acid (50ml) between 5°C and 10°C and the solution was stirred for two hours at 10°C. The  
25 solution was evaporated under reduced pressure and two times coevaporated with toluene. 400ml of ethanol was added and the mixture was stirred for 30 minutes at 40°C. The mixture was cooled and the biproduct filtered. The solution was evaporated under reduced pressure and the product was isolated by silica gel column chromatography.

- 30 Yield: 33g . The intermediate may be esterified to the phenolic hydroxy group of the NNRTI using conventional esterification (ie a coupling agent such as DCC/DMAP but is preferably used as an intermediate linking moiety in conjunction with an methyloxy moiety as described above.



<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) 0.88 (m, 12H) 2.04 (m, 2H) 2.46 (m, 4H) 3.94-4.40 (m, 6H) 5.02 (s, 4H) 5.18 (m, 1H) 7.32 (m, 10H) 7.74 (d, 2H)

Example AA-I-7

5 Alternative route to succinic acid 1,3-bis-(N-CBZ-L-valyloxy)-2-propyl ester

a) Succinic acid 1,3-dibromo-2-propyl ester, 1,1-dimethylethyl ester.

To a solution of 1,3-dibromopropan-2-ol (10.9 g 50 mmole), succinic acid 1,1-dimethylethyl ester (J. Org.Chem 59 (1994) 4864) (10.45g, 60 mmole) and DMAP (0.61 g, 5 mmole) in dichloromethane (180ml) was added DCC (12.4 g, 60 mmole) in portions at about 10°C. The mixture was stirred overnight at room temperature and cooled to about 5°C. The mixture was filtered and the solution was evaporated under reduced pressure. 250ml ethyl acetate was added and the organic phase was washed twice with 5% citric acid, 5% sodium hydrogen carbonate and water. The solution was dried with sodium sulfate and evaporated under reduced pressure. The product was distilled in vacuo. (bp 0,5 135-140°C ) Yield: 16.8 g

b) Succinic acid 1,3-bis-(N-CBZ-L-valyloxy)-2-propyl ester, 1,1-dimethylethyl ester.

To a solution of N-CBZ-L-valine (18.85 g, 75 mmole) in dried DMF (100ml) was added potassium tert.-butoxide (7.85 g, 70 mmole) and the mixture was stirred for one hour at room temperature. A solution of succinic acid 1,3-dibromo-2-propyl ester, 1,1-dimethylethyl ester (9.35g, 25 mmole) in dried DMF (20ml) was added and the mixture was stirred for eighteen hours at 60°C. The potassium bromide was filtered and the solution evaporated under reduced pressure. 300ml of ethyl acetate were added and the organic phase washed twice with 5% sodium hydrogen carbonate and with water. The organic phase was dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography. Yield: 14g

30 c) 1,3-bis-( N-CBZ-L-valyloxy )-2-propyl succinate monoester .

To a cooled solution of succinic acid 1,3-bis-(N-CBZ-L-valyloxy)-2-propyl ester, 1,1-dimethylethyl ester (13 g, 18.18 mmole) in dichloromethane (100ml) was added trifluoroacetic acid (20ml) and the solution was stirred for six hours at room temperature. The solution was evaporated under reduced pressure.

200ml ethyl acetate was added and the organic phase was washed with 5% sodium hydrogen carbonate and water. The solution was evaporated under reduced pressure. Yield: 11.7g . The intermediate may be esterified to the phenolic hydroxy group of the NNRTI using conventional esterification (ie a coupling agent such as DCC/DMAP but is preferably used as an intermediate linking moiety in conjunction with an methyloxy moiety as described above.

#### Example AA-1-8

##### 3-benzyloxycarbonylpropionic acid chloromethyl ester

10

##### a) Succinic acid monobenzyl ester

Succinic anhydride (30 g, 300 mmole) was dissolved in methylene chloride (300 ml). To the solution were added benzyl alcohol (10.2 ml, 100 mmole), 4-dimethylaminopyridine (1.22 g, 10 mmole) and pyridine (48 ml). After 3 hours the reaction mixture was poured in to citric acid aqueous solution. The organic phase was concentrated to small volume and sodium hydrogen carbonate and water were added. Then mixture was stirred for 30 min. The aqueous phase was collected, and to it was added citric acid aqueous solution. The product precipitated out, was collected and dried. 15.3 g.

##### 20 b) 3-benzyloxycarbonylpropionic acid chloromethyl ester

Succinic acid monobenzyl ester (4.16 g, 20 mmole) was dissolved in dioxane (20 ml). To the solution was added tetrabutylammonium hydroxide aqueous solution (40 %, 11.6 ml, 18 mmole). The solution was dried in vacuo and coevaporated with toluene several times. The residue was dissolved in methylene chloride (60 ml) and then chloriodomethane (14.5 ml, 200 mmole) was added to the solution. The reaction solution was stirred for 18 hr and then evaporated and the product was isolated with silica gel column chromatography. 3.64 g

##### c) 3-Benzyloxycarbonylpropionic acid iodomethyl ester.


30 3-Benzyloxycarbonylpropionic acid chloromethyl ester (2 g, 1.38 mmole) was dissolved in acetonitrile (30 ml). Sodium iodide (1.6 g, 10.9 mmole) was added to the solution. After reaction at 70° C for 3 hr, the reaction mixture was filtered and the residue was dissolved in methylene chloride (20 ml) and

refiltered. The solution was dried and gave intermediate 3-benzyloxycarbonylpropionic acid iodomethyl ester in quantitative yield. This intermediate is bonded to an accessible function of a drug, such as a ring hydroxy or carboxy function using conventional alkylation/acylation conditions as described generally herein. Following deprotection of the terminal carboxy, a di/trifunctional linker bearing  $R_2$ , such as 1,3-bis- O-(L-valyl)glycerol or iodomethoxy-L-valyl is acylated/alkylated thereon or  $R_2$  amide bonded thereon by conventional techniques as described herein, such as with DCC coupling agent.

10

### Example AA-I-9

#### 1,3-bis(*N*-tert-butoxycarbonyl-L-valyloxy)-2-propyl 1-iodoethyl carbonate



(a) 1,3-bis(*N*-tert-butoxycarbonyl-L-valyloxy)-2-propyl 1-chloroethyl carbonate.

15

To a solution of 1,3-bis(*N*-tert-butoxycarbonyl-L-valyloxy)-2-propanol (0.545 g, 1.11 mmol) in 5 mL dry  $\text{CH}_2\text{Cl}_2$  were added pyridine (540  $\mu\text{L}$ , 6.68 mmol), with cooling and stirring in an ice bath, followed by 1-chloroethyl chloroformate (242  $\mu\text{L}$ , 2.22 mmol). After 1 h, the reaction mixture was diluted with 5 mL  $\text{CH}_2\text{Cl}_2$  and washed with water (5 mL) and brine (5 mL). The organic phase was dried over anhydrous  $\text{Na}_2\text{SO}_4$  and concentrated on a rotavapor, coevaporating several times with toluene. Column chromatography (silica, 4/1 petroleum ether - ethyl acetate) gave the chloride (596 mg, 90%) as a white solid.

(b) 1,3-bis(*N*-tert-butoxycarbonyl-L-valyloxy)-2-propyl 1-iodoethyl carbonate.

25


A mixture of the chloride (596 mg, 1.0 mmol) from step (a) and NaI (684 mg, 4.57 mmol) in 10 ml dry MeCN was refluxed at 80  $^\circ\text{C}$  for 4 h. The reaction

mixture was concentrated under vacuum and then partitioned between 30 mL diethyl ether and 10 mL water. The organic phase was washed with 5% aqueous sodium thiosulfate (2 x 5 mL), and the last aqueous layer was reextracted with ether (5 mL). The organic phases were combined, washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. Flash column chromatography (silica, 4/1 petroleum ether – ethyl acetate) gave a fraction (275 mg) containing 80% iodide, as determined from <sup>1</sup>H NMR, and small amounts of the starting chloride and alkene from the elimination side reaction.

<sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) 0.81-0.85 (m, 6H), 0.88-0.92 (m, 6H), 1.37 (s, 18H), 2.05 (m, 2H), 2.17 (d, 3H, *J* = 6.1 Hz), 4.12-4.46 (m, 6H), 5.00 (d, 2H, *J* = 8.8 Hz), 5.11 (m, 1H), 6.68 and 6.69 (2 sets of q, 1H, *J* = 6.1 Hz).

Example A-I-10

3-(*N*-benzyloxycarbonyl-L-valyloxy)-2,2-dimethylpropyl iodomethyl carbonate



15

(a) 3-(*N*-benzyloxycarbonyl-L-valyloxy)-2,2-dimethyl-1-propanol.

A mixture of *N*-benzyloxycarbonyl-L-valine (2.50 g, 10.0 mmol), 2,2-dimethyl-1,3-propanediol (5.30 g, 50.9 mmol), dicyclohexylcarbodiimide (2.60 g, 12.6 mmol), and 4-dimethylaminopyridine (125 mg, 1.0 mmol) in 100 mL dry CH<sub>2</sub>Cl<sub>2</sub> was stirred for 23 h. The reaction mixture was filtered and washed successively with 50 mL each of water, saturated aqueous NH<sub>4</sub>Cl, saturated aqueous NaHCO<sub>3</sub>, and water. The organic phase was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated. The title compound (2.99 g, 87%) was isolated by flash column chromatography (silica, 2/1 petroleum ether – ethyl acetate) as a white waxy solid.

(b) 3-(*N*-benzyloxycarbonyl-L-valyloxy)-2,2-dimethylpropyl chloromethyl carbonate

Chloromethyl chloroformate (1.50 mL, 16.6 mmol) was added to a solution of the alcohol (2.74 g, 8.12 mmol) from step (a) and pyridine (4.9 mL, 61 mmol) in 40 mL dry CH<sub>2</sub>Cl<sub>2</sub>, in an ice bath. After stirring for 1 h, the mixture was

30

diluted with  $\text{CH}_2\text{Cl}_2$  and washed successively with water, saturated  $\text{NaHCO}_3$ , and brine. The organic phase was dried over anhydrous  $\text{Na}_2\text{SO}_4$  and concentrated, coevaporating several times with toluene on a rotavapor. Flash column chromatography (silica, 2/1 petroleum ether – ethyl acetate) gave 3.31 g (95%) of the title compound.


c) 3-(*N*-benzyloxycarbonyl-L-valyloxy)-2,2-dimethylpropyl iodomethyl carbonate

A mixture of the chloride (3.14 g, 7.30 mmol) from step (b) and NaI (4.37 g, 29.2 mmol) in 73 mL dry MeCN was refluxed at 80 °C for 3 h. After removal of solvent under vacuum, the mixture was partitioned between 80 mL ethyl acetate and 40 mL water. The organic phase was washed with 5%  $\text{Na}_2\text{S}_2\text{O}_3$ , and then brine, dried over anhydrous  $\text{Na}_2\text{SO}_4$ , and concentrated. Flash column chromatography (silica, petroleum ether – ethyl acetate) gave 3.68 g (97%) of the title compound.

$^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ ) 0.88 and 0.96 (2d, 3H each), 0.98 (s, 6H), 2.18 (m, 1H), 3.94 and 4.02 (2s, 2H each), 4.32 (dd, 1H,  $J = 9.0, 4.7$  Hz), 5.11 (s, 2H), 5.26 (d, 1H), 5.92 and 5.93 (ABq, 2H,  $J_{AB} = 5.1$  Hz), 7.35 (s, 5H).

#### Example AA-I-11

##### 4-(*N*-Boc-L-valyloxy)butyric acid



#### a) Preparation of 4-bromobutyric acid benzyl ester

4-bromobutyric acid (10.6g, 60 mmole) was dissolved in thionyl chloride (20 mmol), and the reaction was kept for 4 hr. The solution was evaporated and coevaporated with toluene several times. The residue was redissolved in dichloromethane (120 ml), and then benzyl alcohol (4.14 ml, 40 mmole) was added. The solution was cooled down to -50° C and triethylamine (10 ml, 72

mmole) was added. The reaction mixture was slowly warmed to room temperature. After 3 hr, the reaction mixture was poured into sodium bicarbonate aqueous solution and the organic phase was washed with water and dried, giving the titled product, 6.8 g.

- 5 b) Preparation of 4-(N-Boc-L-valyloxy)butyric acid benzyl ester  
N-Boc-L-valine (1.3 g, 6 mmole) was dissolved in dioxane (5 ml). To the solution was added tetrabutylammonium hydroxide aqueous solution (40 %, 3.8 ml, 6 mmole), and the solution was evaporated and coevaporated with toluene several times. The residue was dissolved in DMF (15 ml) and 4-  
10 bromobutyric acid benzyl ester (1.28g, 5mmole) was added to it. The reaction was kept for 18 hr, and then poured into sodium bicarbonate aqueous solution and extracted with dichloromethane. The organic phase was dried and the product was isolated with silica gel column chromatography, 1.2 g.


c) 4-( N-Boc-L-valyloxy)butyric acid.

- 15 To a solution of 4-(N-Boc-L-valyloxy)butyric acid benzyl ester (1.2 g , 3 mmole) in ethyl acetate/methanol (5ml/5ml) was added palladium black (20 mg). The reaction mixture was kept under hydrogen at atmospheric pressure for 2 hr. The suspension was filtered through Celite and dried, giving the title product, 840 mg. The intermediate may be esterified to the phenolic hydroxy  
20 group of the NNRTI using conventional esterification (ie a coupling agent such as DCC/DMAP but is preferably used as an intermediate linking moiety in conjunction with an methyloxy moiety as described above.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 5.05 (d, 1H) 4.20 (m, 3H) 2.48 (t, 2H) 2.00 (m, 2H) 1.46 (s, 9H) 0.96 (m, 6H).

#### Example AA-I-12

##### N-BOC-L-isoleucine iodomethyl ester



- 30 a) N-BOC-L-isoleucine chloromethyl ester.

To a solution of N-BOC-L-isoleucine (23.1 g, 0.1 mol) in dioxane (500 mL), was added dropwise a 40% aqueous solution of tetrabutylammonium hydroxide (65.6 mL, 0.1 mol). After stirring for 15 min, the solution was evaporated to dryness through co-evaporation with dioxane and toluene. The residue was dissolved in dichloromethane (500 mL) and then chloriodomethane (72.8 mL, 1 mol) was added and the solution was stirred for 6h at room temperature. The solution was concentrated under reduced pressure and the residue was shaken with hexane / ethyl acetate (1:1 v/v, 400 mL). The yellow crystalline solid was filtered off and the filtrate was washed with aqueous solution of sodium thiosulfate (0.1 M) and then filtered through anhydrous sodium sulfate and evaporated to dryness. The residue was column chromatographed (silica gel, 1-2% MeOH in CH<sub>2</sub>Cl<sub>2</sub>), to give 20.8 g of N-BOC-L-isoleucine chloromethyl ester.


b) N-BOC-L-isoleucine iodomethyl ester.

15 To a solution of N-BOC-L-isoleucine chloromethyl ester (19.6 g, 70 mmol) in acetonitrile (300 mL), was added sodium iodide (31.5 g, 210 mmol). The solution was stirred for 4 h at 60 °C. The resulting suspension was filtered and the filtrate was evaporated. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and washed with aqueous sodium thiosulfate (0.1 M). The organic phase was dried  
20 (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure. The crude product was column chromatographed (silica gel, 2% MeOH in CH<sub>2</sub>Cl<sub>2</sub>), to give 22.6 g of N-BOC-L-isoleucine iodomethyl ester.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 6.04 (d, 1H), 5.82 (d, 1H), 4.97 (d, 1H), 4.25 (dd, 1H), 1.98-  
25 1.80 (m, 1H), 1.43 (s, 9H), 1.50-1.05 (m, 2H), 0.97-0.88 (m, 6H).

### Example AA-I-13

2,2-dimethyl-3-(N-Boc-L-valyloxy)propionic acid iodomethyl ester



30

a) 2,2-dimethyl-3-( N-Boc-L-valyloxy)propionic acid:

N-Boc-L-valine (10.8g, 50 mmole), 4-dimethylaminopyridine (610 mg, 5 mmole) and DCC (6.18 g, 30 mmole) were dissolved in methylene chloride (100 ml). After stirring for 2 hour the mixture was filtered. To the filtrate were added 2,2-dimethyl-3-hydroxy-propionic acid (3.54g, 30 mmole) and pyridine (10 ml). After 18 hr, the reaction mixture was filtered, and the filtrate was poured into sodium hydrogen carbonate aqueous solution, the organic phase was then washed with citric acid aqueous solution and water succesively. After evaporation the product was isolated with silica gel column chromatography to yield 4.4g.

b) 2,2-dimethyl-3-(N-Boc-L-valyloxy)propionic acid chloromethyl ester . 2,2-dimethyl-3-(N-Boc-L-valyloxy)propionic acid (3.9 g, 12.3 mmole) was dissolved in dioxane (60 ml). To the solution was added tetrabutylammonium hydroxide aqueous solution (40 %, 7.78 ml, 12 mmole). The solution was dried in vacuo, and it was coevaporated with toluene for several times. The residue was dissolved in methylene chloride and then chloriodomethane (18.9 ml, 260 mmole) was added to the solution. After 18 hr, the reaction solution was evaporated and the product was isolated with silica gel column chromatography to yield 3.7 g.

c) 2,2-dimethyl-3-(N-Boc-L-valyloxy)propionic acid iodomethyl ester . 2,2-Dimethyl-3-( N-Boc-L-valyloxy)propionic acid chloromethyl ester ( 3.6 g, 10 mmole ) was dissolved in acetonitrile (50 ml). Sodium iodide (2.1 g, 14 mmole) was added to the solution. After reaction at 70° C for 2 hr, the reaction mixture was filtered and the residue was dissolved in methylene chloride (20 ml) and refiltered. The solution was dried and gave 4.34g of the titled product.


25

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 5.92 (dd, 2H) 5.10 (d, 1H) 4.24 (m, 1H) 4.15 (dd, 2H) 2.01 (m, 1H) 1.44 (s, 9H) 1.25 (d, 6H) 0.91 (m, 6H )

#### Example AA-I-14

30 3,3- bis (N-CBz-L-valyloxymethyl)-propionic acid iodomethyl ester





a) Preparation of 3,3-bis (N-CBz-L-valyloxymethyl)-propionic acid chloromethyl ester.

3,3- bis (N-CBz-L-valyloxymethyl)-propionic acid (3 g, 5 mmole) was dissolved  
 5 in dioxane (20 ml). To the solution was added tetrabutylammonium hydroxide  
 aqueous solution (40 %, 3.11 ml, 4.8 mmole). The solution was dried in  
 vacuo, and it was coevaporated with toluene several times. The residue was  
 dissolved in methylene chloride (15 ml) and then chloriodomethane (7.3 ml,  
 100 mmole) was added to the solution. The reaction solution was refluxed for  
 10 18 hr and then evaporated and the product was isolated with silica gel column  
 chromatography. 900 mg.


b) 3,3-bis-(N-CBz-L-valyloxymethyl)propionic acid iodomethyl ester..

3,3-bis (N-CBz-L-valyloxymethyl)-propionic acid chloromethyl ester (900 mg,  
 1.38 mmole) was dissolved in acetonitrile (5 ml). Sodium iodide (289 mg, 1.93  
 15 mmole) was added to the solution. After reaction at 70° C for 3 hr, the reaction  
 mixture was filtered and the residue was dissolved in methylene chloride (5  
 ml) and refiltered. The solution was dried and gave the titled product. 800 mg.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 7.35 (m, 10 H) 5.88 (dd, 2H) 5.25 (d, 2H) 4.29 (m, 2H) 4.18  
 20 (m, 4H) 2.56 (m, 1H) 2.42 (d, 2H) 2.16 (m, 2H) 0.93 (m 12 H)

#### Example AA-I-15

##### 2-(N-CBz-L-valyloxy)ethoxycarbonyloxymethyl iodide




2-(N-CBz-L-valyloxy)ethoxycarbonyloxymethyl chloride (1.16 g, 3 mmole) was dissolved in acetonitrile (10 ml). Sodium iodide (630 g, 4.2 mmole) was added to the solution. After reaction at 65° C for 2.5 hr, the reaction mixture was cooled down to room temperature and filtered and the residue was dissolved in methylene chloride (5 ml) and refiltered. The solution was dried and gave the titled product. 1.2 g.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 7.35 (m, 5H) 5.93 (dd, 2H) 5.26 (d, 1H) 5.11 (s, 2H) 4.39 (m, 5H) 2.18 (m, 1H) 0.94 (m, 6 H).

#### Example AA-I-16

##### 1,3-bis(N-tert-butoxycarbonyl-L-valyloxy)-2-propyl iodomethyl carbonate)



##### a) 1-O-(N-tert-butoxycarbonyl-L-valyl)glycerol

*N*-tert-Butoxycarbonyl-L-valine (32.53 g, 0.150 mol), *N,N'*-dicyclohexylcarbodiimide (37.85 g, 0.183 mol, and 4-dimethylaminopyridine (1.83 g, 0.015 mol) were added to glycerol (138.12 g, 1.5 mol) in 500 mL dry DMF and the mixture was stirred at rt under N<sub>2</sub> for 3 days. The reaction mixture was filtered, concentrated under vacuum, and then partitioned between 300 mL EtOAc and 150 mL H<sub>2</sub>O. The aqueous phase was reextracted with 150 mL EtOAc. The organic phases were combined and washed successively with 100 mL each of saturated aqueous NaHCO<sub>3</sub>, saturated NH<sub>4</sub>Cl, and brine. Drying over anhydrous Na<sub>2</sub>SO<sub>4</sub>, and concentration under vacuum gave a viscous light yellow oil as crude product. Flash column chromatography on silica gel with 4/1 EtOAc - petroleum ether (BP 40-60 °C) gave 18.27 g (42%) of product (alternative nomenclature: 3-(N-tert-butoxycarbonyl-L-valyloxy)-1,2-propanediol). Reactions done overnight gave similar yields.

##### b) 1,3-di-O-(N-tert-butoxycarbonyl-L-valyl)glycerol

1-O-(N-tert-butoxycarbonyl-L-valyl)glycerol (17.95 g, 61.6 mmol), Boc-L-valine (6.69 g, 30.8 mmol), DMAP (0.38 g, 3.1 mmol), and DCC (7.10 g, 34.4 mmol) in 240 mL CH<sub>2</sub>Cl<sub>2</sub> and 60 mL DMF were stirred at rt under N<sub>2</sub> for 18 h. The reaction mixture was filtered, concentrated under vacuum, and redissolved in

200 mL EtOAc. The organic solution was washed with 50 mL saturated  $\text{NH}_4\text{Cl}$ . The aqueous phase was reextracted with 50 mL EtOAc. The organic phases were combined, washed successively with 50 mL saturated  $\text{NaHCO}_3$  and 50 mL brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under vacuum. Flash column chromatography of the crude material on silica gel (eluent 2/1 petroleum ether – EtOAc, and then EtOAc) gave 7.41 g (49%) of the title compound (alternative nomenclature: 1,3-bis(*N*-tert-butoxycarbonyl-L-valyloxy)-2-propanol).

c) 2-O-chloromethoxycarbonyl-1,3-di-O-(*N*-tert-butoxycarbonyl-L-valyl)glycerol.

Chloromethyl chloroformate (2.70 mL, 30 mmol) was added to a solution of 1,3-di-O-(*N*-tert-butoxycarbonyl-L-valyl)glycerol (7.27 g, 14.8 mmol) and pyridine (7.2 mL, 89 mmol) in 60 mL dry  $\text{CH}_2\text{Cl}_2$ , in an ice bath, under  $\text{N}_2$ . After stirring for 1 h 45 min, the reaction mixture was diluted with 100 mL  $\text{CH}_2\text{Cl}_2$  and washed with 40 mL water. The aqueous phase was reextracted with 20 mL  $\text{H}_2\text{O}$ . The organic phases were combined, washed with 40 mL saturated  $\text{NaHCO}_3$ , followed by 2 x 50 mL brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated under vacuum. Flash column chromatography on silica gel with 2/1 hexane- EtOAc gave 8.03 g (93%) of the title compound (alternative nomenclature: 1,3-bis(*N*-tert-butoxycarbonyl-L-valyloxy)-2-propyl chloromethyl carbonate).

d) 2-O-iodomethoxycarbonyl-1,3-di-O-(*N*-tert-butoxycarbonyl-L-valyl)glycerol.

A solution of 2-O-chloromethoxycarbonyl-1,3-di-O-(*N*-tert-butoxycarbonyl-L-valyl)propane-1,2,3-triol (7.86 g, 13.5 mmol) and NaI (8.09 g, 54.0 mmol) in 135 mL dry acetonitrile was refluxed at 80 °C for 4 h under  $\text{N}_2$ . The reaction mixture was concentrated under vacuum, and then partitioned between 150 mL diethyl ether and 50 mL  $\text{H}_2\text{O}$ . The aqueous layer was reextracted with 2 x 25 mL ether. The combined organic phases were washed successively with 25 mL aqueous  $\text{Na}_2\text{S}_2\text{O}_3$  and 50 mL brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated. Flash column chromatography (silica gel, 2/1 hexane-ethyl acetate gave 8.38 g (92%) title product (alternative name: 2-iodomethoxycarbonyloxy-1,3-bis-(*N*-tert-butoxycarbonyl-L-valyloxy)propane or

1,3-bis(N-tert-butoxycarbonyl-L-valyloxy)-2-propyl iodomethyl carbonate) as a white solid.

<sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) 0.81 (d, 6H), 0.88 (m, 6H), 1.36 (s, 18H), 2.06 (m, 2H), 4.11-4.46 (m, 6H), 5.0 (br d, 2H), 5.12 (m, 1H), 5.88 (s, 2H).

#### Example A-I-1

##### Iodomethyl 2-methyl-2-(N-benzyloxycarbonyl-L-valyloxymethyl) propionate

10 a) 4-Methoxybenzyl 2-(hydroxymethyl)-2-methyl propionate.  
2-(Hydroxymethyl)-2-methyl propionic acid was esterified by alkylation with 4-methoxybenzyl chloride by conventional means, namely treatment with aqueous NaOH, followed by evaporation and dissolution in an organic solvent such as DMF to which the 4-methoxybenzyl chloride is added and the reaction  
15 warmed and agitated, such as stirring at 60 C for one hour. The reaction mixture is cooled, concentrated by rotavapor and the resulting concentrated suspension partitioned between water and dichloromethane. The organic phase is evaporated and the residue subjected to silica gel column chromatography, for example with 0, 2, 4% EtOH in dichloromethane to yield  
20 the title compound (7.10 g). R<sub>f</sub> (2%MeOH/CHCl<sub>3</sub>) 0.40.

b) 4-Methoxybenzyl 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-methyl propionate.  
4-Methoxybenzyl 2-(hydroxymethyl)-2-methyl propionate (2.50 g, 10.5 mmol), N-benzyloxy carbonyl-L-valine (2.51 g, 10 mmole), 4-dimethylaminopyridine (183 mg) and 1-hydroxybenzotriazole (1.35g, 10 mmole) were mixed and dissolved in N,N-dimethylformamide (90 ml). Then dicyclohexyl-carbodiimide (2.47 g 12 mmol) was added. After stirring for 3 days at room temperature the suspension was filtered and the filtrate evaporated in vacuo. The residue was partitioned between 0.1M citric acid and dichloromethane. The organic phase  
25 was then extracted with aqueous saturated NaHCO<sub>3</sub> and evaporated in vacuo. The residue was silica gel column chromatographed (0, 1, 2, 3% ethanol in dichloromethane). The appropriate fractions were pooled and evaporated in vacuo to give the title compound (2.72 g).

d) 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-methyl propionic acid.

To a solution of 4-methoxybenzyl 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-methyl propionate (2.72 g, 5.76 mmole), was added trifluoroacetic acid (11.5 ml) and the emerging dark red solution was stirred for 30 min at room temperature. The solution was evaporated to dryness with dioxane and toluene. The residue was silica gel column chromatographed (2, 3, 4% ethanol in dichloromethane). The appropriate fractions were pooled and evaporated in vacuo to give the title compound (1.86 g).

R<sub>f</sub> (2%MeOH/CHCl<sub>3</sub>) 0.30.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 7.32 (s, 5H), 5.32 (d, 1H), 5.10 (s, 2H), 4.32 (d,d, 1H), 4.21 (d,d, 2H), 2.13 (m, 1H), 1.26 (s, 3H), 1.25 (s, 3H), 0.95 (d, 3H), 0.86 (d, 3H).

c) Chloromethyl 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-methyl propionate.

2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-methyl propionic acid was esterified by conventional techniques, namely dissolution in an organic solvent such as dioxane and dropwise addition of aqueous tetrabutylammonium hydroxide, followed by evaporation. The residue is dissolved in dichloromethane and then chloriodomethane and the mixture stirred for 6 hours at room temperature, followed by partition, shaking the filtrate with aqueous sodium thiosulphate. 0.1M, filtration and evaporation. The title compound (1.40 g) was obtained after silica gel column chromatography (0, 1, 2, 3% ethanol in dichloromethane).

d) Iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-methyl propionate.


Chloromethyl 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-methyl propionate was converted to iodide by conventional techniques, namely addition to NaI in acetonitrile, stirring and heating, for instance to 75 C for four hours. The resulting suspension is filtered and the filtrate evaporated, dissolved in organic solvent such as toluene and shaken with aqueous sodium thiosulphate (0.1M) and evaporation to give the title compound (1.25 g) practically pure.

R<sub>f</sub> (2%MeOH/CHCl<sub>3</sub>) 0.80.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 7.35 (s, 5H), 5.90 (d,d, 2H), 5.24 (d, 1H), 5.10 (s, 2H), 4.31 (d,d, 1H), 4.14 (d,d, 2H), 2.16 (m, 1H), 1.22 (s, 6H), 0.96 (d, 3H), 0.87 (d, 3H).

#### Example A-I-2


- 5 Iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxy)-DL-propionate.



- a) Chloromethyl 2-(N-benzyloxycarbonyl-L-valyloxy)-DL-propionate.  
2-(N-benzyloxycarbonyl-L-valyloxy) propionic acid (1 g) was esterified by the method described in Example A-I-I, step d. The title compound (0.76 g) was  
10 obtained after silica gel column chromatography (0, 1% ethanol in dichloromethane).  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.75.
- b) Iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxy)-DL-propionate  
Chloromethyl 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-methyl propionate was converted to iodide by the method described in Example A-I-1, step e to  
15 give the title compound (0.95 g) practically pure.  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.75.  
 $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 7.33 (s, 5H), 5.98 (d, 1H), 5.86 (d, 1H), 5.26 (d, 1H), 5.10 (s, 2H), 5.07 (q, 1H), 4.38 (d,d, 1H), 2.30 (m, 1H), 1.49 (d, 3H), 1.03 (d, 3H), 0.95 (d, 3H).

#### 20 Example A-I-3

Iodomethyl (1,3-di-(N-benzyloxycarbonyl)-L-valyloxy)-2-propyl carbonate.



- 25 a) Chloromethyl (1,3-di-(N-benzyloxycarbonyl)-L-valyloxy)-2-propyl carbonate.

To a solution of 1,3-di-((N-benzyloxycarbonyl)-L-valyloxy)propan-2-ol (1.34 g, 2.4 mmole) in dichloromethane (10 ml) was added dry pyridine (1.15 ml, 14.4 mmol) and chloromethyl chloroformate (0.43 ml, 4.8 mmole) at  $0^\circ\text{C}$ . The

reaction was then stirred for 30 min and then poured into aqueous 50% saturated sodium chloride / 0.1M citric acid solution and extracted with dichloromethane. The organic phase was evaporated and the residue silica gel column chromatographed (0, 1, 1.5% ethanol in dichloromethane). The appropriate fractions were pooled and evaporated in vacuo to give the title compound (1.26g).  $R_f$  (2%MeOH/  $\text{CHCl}_3$ ) 0.85.

b) Preparation of iodomethyl (1,3-di-(N-benzyloxycarbonyl)-L-valyloxy)-2-propyl carbonate.


Chloromethyl (1,3-di-(N-benzyloxycarbonyl)valyloxy)-2-propyl carbonate was converted to iodide by the method described in Example A-I-1, step e) to give the title compound (1.37 g) practically pure.

$R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.85.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 7.34 (s, 10H), 5.93 (d, 1H), 5.89 (d, 1H), 5.21 (m, 3H), 5.11 (s, 4H), 4.50-4.17 (m, 6H), 2.12 (m, 2H), 0.97 (d, 6H), 0.88 (d, 6H).

#### Example A-I-4

##### Iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxy)isobutyrate.



a) 4-Methoxybenzyl 2-hydroxyisobutyrate.

2-hydroxy isobutyric acid (1.56 g) was esterified by alkylation with 4-methoxybenzyl chloride by the method described in Example A-I-1, step a).

The title compound (2.65 g) was obtained after silica gel column chromatography (0, 1, 2% ethanol in dichloromethane).  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.45.

b) 4-Methoxybenzyl 2-(N-benzyloxycarbonyl-L-valyloxy) isobutyrate.

4-methoxybenzyl 2-hydroxyisobutyrate was acylated with N-benzyloxycarbonyl-L-valine by the method described in Example A-I-1, step b). The title compound (3.21 g) was obtained after silica gel column

chromatography (0, 1, 1.5% ethanol in dichloromethane).  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.70.

c) 2-(N-benzyloxycarbonyl-L-valyloxy) isobutyric acid.

4-methoxybenzyl 2-(N-benzyloxycarbonyl-L-valyloxy) isobutyrate was de-  
esterified by the method described in Example A-I-1 step c. The title  
compound (2.01 g) was obtained after silica gel column chromatography (2,  
5 10, 20% ethanol in dichloromethane).  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.30.  
 $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 7.32 (s, 5H), 5.33 (d, 1H), 5.10 (s, 2H), 4.31 (d,d, 1H), 2.22  
(m, 1H), 1.57 (s, 6H), 0.98 (d, 3H), 0.89 (d, 3H).

d) Chloromethyl 2-(N-benzyloxycarbonyl-L-valyloxy) isobutyrate.

2-(N-benzyloxycarbonyl-L-valyloxy) isobutyric acid was esterified by the  
10 method described in Example A-I-1, step d. The title compound (1.90 g) was  
obtained after silica gel column chromatography (0, 1, 1.5% ethanol in  
dichloromethane).  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.80.

e) Iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxy) isobutyrate.


Chloromethyl 2-(N-benzyloxycarbonyl-L-valyloxy) isobutyrate was converted  
15 to iodide by the method described in Example A-I-1, step e to give the title  
compound (2.32 g) practically pure.

$R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.80.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 7.33 (s, 5H), 5.89 (s, 2H), 5.22 (d, 1H), 5.11 (s, 2H), 4.29  
20 (d,d, 1H), 2.21 (m, 1H), 1.55 (s, 3H), 1.53 (s, 3H), 1.00 (d, 3H), 0.93 (d, 3H).

#### EXAMPLE A-I-5

Iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxy)-3-methyl-(S)-(+)-butyrate.



25 a) 4-Methoxybenzyl 2-hydroxy-3-methyl-(S)-(+)-butyrate.

2-hydroxy-3-methyl-(S)-(+)-butyric acid (1.77 g) was esterified by alkylation  
with 4-methoxybenzyl chloride by the method described in Example A-I-1,  
step a. The title compound (3.10 g) was obtained after silica gel column  
chromatography (0, 1, 2% ethanol in dichloromethane).  $R_f$  (2%MeOH/ $\text{CHCl}_3$ )  
30 0.50.



b) 4-Methoxybenzyl 2-(N-benzyloxycarbonyl-L-valyloxy)-3-methyl-(S)-(+)-butyrate.

4-Methoxybenzyl 2-hydroxy-3-methyl-(S)-(+)-butyrate was acylated with N-benzyloxycarbonyl-L-valine by the method described in Example A-I-1, step b.

5 The title compound (5.74 g) was obtained after silica gel column chromatography (0, 1, 1.5% ethanol in dichloromethane).  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.70.

c) 2-(N-benzyloxycarbonyl-L-valyloxy)-3-methyl-(S)-(+)-butyric acid.

4-methoxybenzyl 2-(N-benzyloxycarbonyl-L-valyloxy)-3-methyl-(S)-(+)-

10 butyrate was de-esterified by the method described in Example A-I-1, step c.

The title compound (3.41 g) was obtained after silica gel column

chromatography (2, 10, 20% ethanol in dichloromethane).  $R_f$

(2%MeOH/ $\text{CHCl}_3$ ) 0.45.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 7.36 (s, 5H), 5.38 (d, 1H), 5.11 (s, 4H), 4.90 (d, 1H), 4.41 (d,d, 1H), 2.28 (m, 2H), 1.04-0.89 (m, 12H).

15 d) Chloromethyl 2-(N-benzyloxycarbonyl-L-valyloxy)-3-methyl-(S)-(+)-butyrate.

2-(N-benzyloxycarbonyl-L-valyloxy)-3-methyl-(S)-(+)-butyric acid was

esterified by the method described in Example A-I-1, step d. The title

compound (2.96 g) was obtained after silica gel column chromatography (0, 1,

20 2% ethanol in dichloromethane).  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.85.

e) Iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxy)-3-methyl-(S)-(+)-butyrate.

Chloromethyl 2-(N-benzyloxycarbonyl-L-valyloxy)-3-methyl-(S)-(+)-butyrate


was converted to iodide by the method described in Example A-I-1, step e to

25 give the title compound (3.64 g) practically pure.

$R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.85.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 7.36 (s, 5H), 6.00 (d, 1H), 5.83 (d, 1H), 5.28 (d, 1H), 5.11 (s, 4H), 4.83 (d, 1H), 4.41 (d,d, 1H), 2.29 (m, 2H), 1.05-0.90 (m, 12H).

## EXAMPLE A-I-6

Iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxy)-2-phenyl-DL-acetate.


- 5 a) 4-Methoxybenzyl 2-hydroxy-2-phenyl-DL-acetate.  
DL-mandelic acid (2.28 g) was esterified by alkylation with 4-methoxybenzyl chloride by the method described in Example A-I-1, step a. The title compound (3.69 g) was obtained after silica gel column chromatography (0, 1, 1.5% ethanol in dichloromethane).  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.55.
- 10 b) 4-Methoxybenzyl 2-(N-benzyloxycarbonyl-L-valyloxy)-2-phenyl-DL-acetate.  
4-Methoxybenzyl 2-hydroxy-2-phenyl-DL-acetate was acylated with N-benzyloxycarbonyl-L-valine by the method described in Example A-I-1, step b. The title compound (6.50 g) was obtained after silica gel column
- 15 chromatography (0, 1, 1.5% ethanol in dichloromethane).  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.75.
- c) 2-(N-benzyloxycarbonyl-L-valyloxy)-2-phenyl-DL-acetic acid.  
4-Methoxybenzyl 2-(N-benzyloxycarbonyl-L-valyloxy)-2-phenyl-DL-acetate was de-esterified by the method described in Example A-I-1, step c. The title
- 20 compound (4.75 g) was obtained after silica gel column chromatography (2, 10, 20% ethanol in dichloromethane).  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.40.  
 $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 7.36 (m, 10H), 5.91 (d, 1H), 5.27 (m, 1H), 5.04 (s, 2H), 4.57-4.40 (2xd,d, 1H), 2.30 (m, 1H), 1.01-0.82 (m, 6H).
- d) Chloromethyl 2-(N-benzyloxycarbonyl-L-valyloxy)-2-phenyl-DL-acetate.  
25 2-(N-benzyloxycarbonyl-L-valyloxy)-2-phenyl-DL-acetic acid was esterified by the method described in Example A-I-1, step d. The title compound (1.69 g) was obtained after silica gel column chromatography (0, 1, 2% ethanol in dichloromethane).  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.80.
- e) Iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxy)-2-phenyl-DL-acetate.

Chloromethyl 2-(N-benzyloxycarbonyl-L-valyloxy)-2-phenyl-DL-acetate was converted to iodide by the method described in Example A-I-1, step e to give the title compound (1.89 g) practically pure.

- 5  $R_f$  (2%MeOH/CHCl<sub>3</sub>) 0.80.  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 7.36 (m, 10H), 5.94-5.82 (m, 3H), 5.28 (m, 1H), 5.10 (s, 2H), 4.46 (m, 1H), 2.21 (m, 1H), 1.08-0.85 (m, 6H).

#### Example A-I-7

- 10 Iodomethyl 4-(N-benzyloxycarbonyl-L-valyloxy) benzoate.



- a) 4-Methoxybenzyl 4-hydroxybenzoate.  
4-Hydroxybenzoic acid (1.38 g) was esterified by alkylation with 4-methoxybenzyl chloride by the method described in Example A-I-1, step a.
- 15 The title compound (2.06 g) was obtained after silica gel column chromatography (0, 1, 2, 3% ethanol in dichloromethane).  $R_f$  (2%MeOH/CHCl<sub>3</sub>) 0.40.
- b) 4-Methoxybenzyl 4-(N-benzyloxycarbonyl-L-valyloxy) benzoate.  
4-Methoxybenzyl 4-hydroxybenzoate was acylated with N-benzyloxycarbonyl-L-valine by the method described in Example A-I-1, step b. The title compound (2.71 g) was obtained after silica gel column chromatography (0, 1% ethanol in dichloromethane).  $R_f$  (2%MeOH/CHCl<sub>3</sub>) 0.70.
- 20 c) 4-(N-benzyloxycarbonyl-L-valyloxy) benzoic acid.  
4-Methoxybenzyl 4-(N-benzyloxycarbonyl-L-valyloxy) benzoate was de-esterified by the method described in Example A-I-1, step c. The title compound (1.86 g) was obtained after silica gel column chromatography (2, 10, 20% ethanol in dichloromethane).  $R_f$  (2%MeOH/CHCl<sub>3</sub>) 0.20.  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 8.15 (d, 2H), 7.34 (m, 5H), 7.22 (d, 2H), 5.38 (d, 1H), 5.17 (s, 2H), 4.58 (d,d, 1H), 2.34 (m, 1H), 1.12 (s, 3H), 0.96 (d, 3H).
- 30 d) Chloromethyl 4-(N-benzyloxycarbonyl-L-valyloxy) benzoate.

4-(N-benzyloxycarbonyl-L-valyloxy) benzoic acid was esterified by the method described in Example A-I-1, step d. The title compound (0.95 g) was obtained after silica gel column chromatography (0, 1% ethanol in dichloromethane).  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.80.


- 5 e) Iodomethyl 4-(N-benzyloxycarbonyl-L-valyloxy) benzoate.  
Chloromethyl 4-(N-benzyloxycarbonyl-L-valyloxy) benzoate was converted to iodide by the method described in Example A-I-1, step e to give the title compound (1.16 g) practically pure.

10  $R_f$  (2%MeOH/ $\text{CHCl}_3$ ) 0.80.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 8.11 (d, 2H), 7.35 (m, 5H), 7.21 (d, 2H), 6.15 (s, 2H), 5.32 (d, 1H), 5.14 (s, 2H), 4.55 (d,d, 1H), 2.34 (m, 1H), 1.10 (s, 3H), 1.03 (d, 3H).

#### Example A-1-8

- 15 Iodomethyl 5-(N-CBz-L-valyloxy)-2,2-dimethylvalerate



- a) 4-Methoxybenzyl 2,2-dimethyl-4-pentenoate

To a solution of 2,2-dimethyl-4-pentenoic acid (11.5 g, 90 mmol) in DMF (250 mL) at room temperature, was added potassium tert-butoxide (11.1 g, 99 mmol). The reaction mixture was stirred at 60 °C for 1h. 4-Methoxybenzyl chloride (16.9 g, 108 mmol) was added and the reaction mixture was stirred at 60 °C for 4h. The DMF was evaporated under vacuum, the residue was dissolved in ether (500 mL) and washed with water (3 x 200 mL). The organic phase was dried with  $\text{Na}_2\text{SO}_4$  and evaporated to give 21.4 g of 4-methoxybenzyl 2,2-dimethyl-4-pentenoate.

- b) 4-Methoxybenzyl 5-hydroxy-2,2-dimethylvalerate

A mixture of 4-methoxybenzyl 2,2-dimethyl-4-pentenoate (9.50 g, 38 mmol) and 9-BBN (115 mL, 57 mmol, 0.5 M in THF) was stirred at 60 °C for 60 min, whereupon the reaction mixture was cooled to -5 °C.  $\text{H}_2\text{O}$  (35 mL) was added, the reaction mixture was stirred for 5 min at -5 °C, an aqueous solution of NaOH (35 mL, 3M) was added and the reaction mixture was stirred for a

further 10 min at -5 °C. An aqueous solution of H<sub>2</sub>O<sub>2</sub> (35 mL, 30%) was added dropwise and the temperature of the reaction mixture was allowed to assume room temperature, whereupon the reaction mixture was stirred for 30 min at room temperature. After evaporation, water (200 mL) was added and the  
5 resulting mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (5 x 200 mL). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure. The crude product was column chromatographed (silica gel, 1→8% MeOH in CH<sub>2</sub>Cl<sub>2</sub>), to give 6.32 g of 4-methoxybenzyl 5-hydroxy-2,2-dimethylvalerate.

10 c) 4-Methoxybenzyl 5-(N-CBz-L-valyloxy)-2,2-dimethylvalerate

To a mixture of DCC (9.41 g, 46 mmol), DMAP (0.586 g, 4.8 mmol) and N-CBz-L-valine (12.1 g, 48 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (200 mL) at 0 °C, was added dropwise a solution of 4-methoxybenzyl 5-hydroxy-2,2-dimethyl-valerate (6.40 g, 24 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (50 mL). After 1h at 0 °C, the temperature of the  
15 reaction mixture was allowed to assume room temperature and then the mixture was stirred for 5h at room temperature. The mixture was filtered through a glass filter and the solvent was removed under reduced pressure. The crude product was column chromatographed (silica gel, 1→4% MeOH in CH<sub>2</sub>Cl<sub>2</sub>), to give 8.61 g 4-methoxybenzyl 5-(N-CBz-L-valyloxy)-2,2-dimethylvalerate.  
20

d) 5-(N-CBz-L-valyloxy)-2,2-dimethylvaleric acid

To a solution of 4-methoxybenzyl 5-(N-CBz-L-valyloxy)-2,2-dimethylvalerate (8.24 g, 16.5 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (100 mL) at room temperature, was added trifluoroacetic acid (5 mL). After 1h at room temperature, the reaction mixture  
25 was concentrated under reduced pressure. The crude product was column chromatographed (silica gel, 3→5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>), to give 6.00 g of 5-(N-CBz-L-valyloxy)-2,2-dimethylvaleric acid.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 10.94 (br s, 1H), 7.35 (s, 5H), 5.45 (d, 1H), 5.11 (s, 2H), 4.30 (dd, 1H), 4.21-4.00 (m, 2H), 2.28-2.07 (m, 1H), 1.68-1.51 (m, 4H), 1.21  
30 (s, 6H), 0.97 (d, 3H), 0.89 (d, 3H).

e) Chloromethyl 5-(N-CBz-L-valyloxy)-2,2-dimethylvalerate

To a solution of 5-(N-CBz-L-valyloxy)-2,2-dimethylvaleric acid (5.88 g, 15.5 mmol) in dioxane (100 mL), was added dropwise a 40% aqueous solution of

tetrabutylammonium hydroxide (10.1 g). After stirring for 5 min, the solution was evaporated to dryness through co-evaporation with dioxane and toluene. The residue was dissolved in dichloromethane (100 mL) and then chloriodomethane (11.3 mL, 155 mmol) was added and the solution was stirred for 6 h at room temperature. The solution was concentrated under reduced pressure and the residue was shaken with hexane / ethyl acetate (1:1 v/v, 200 mL). The yellow crystalline solid was filtered off and the filtrate was washed with aqueous solution of sodium thiosulfate (0.1 M) and the filtered through anhydrous sodium sulfate and evaporated to dryness. The residue was column chromatographed (silica gel, 1-4% MeOH in CH<sub>2</sub>Cl<sub>2</sub>), to give 3.95 g of chloromethyl 5-(N-CBz-L-valyloxy)-2,2-dimethylvalerate.


f) Iodomethyl 5-(N-CBz-L-valyloxy)-2,2-dimethylvalerate.

To a solution of chloromethyl 5-(N-CBz-L-valyloxy)-2,2-dimethylvalerate (3.85 g, 9 mmol) in acetonitrile (50 mL), was added sodium iodide (5.40 g, 36 mmol). The solution was stirred for 4 h at 60 °C. The resulting suspension was filtered and the filtrate was evaporated. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and washed with aqueous sodium thiosulfate (0.1 M). The organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure. The crude product was column chromatographed (silica gel, 1% MeOH in CH<sub>2</sub>Cl<sub>2</sub>), to give 4.26 g of iodomethyl 5-(N-CBz-L-valyloxy)-2,2-dimethylvalerate.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 7.34 (s, 5H), 5.90 (s, 2H), 5.32 (d, 1H), 5.10 (s, 2H), 4.29 (dd, 1H), 4.18-4.02 (m, 2H), 2.26-2.08 (m, 1H), 1.65-1.50 (m, 4H), 1.17 (s, 6H), 0.97 (d, 3H), 0.89 (d, 3H).

#### Example A-1-9

##### 2-(N-CBz-L-valyloxy)-ethyl iodomethyl carbonate



a) 2-(N-CBz-L-valyloxy)-ethanol:

To a mixture of DCC (11.4 g, 55 mmol), DMAP (0.611 g, 5 mmol) and ethyleneglycol (55.8 mL, 1 mol) in CH<sub>2</sub>Cl<sub>2</sub> (300 mL) at 0 °C, was added dropwise a solution of N-CBz-L-valine (12.6 g, 50 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (100 mL).

After 1h at 0 °C, the temperature of the reaction mixture was allowed to assume room temperature and then the mixture was stirred for 5h at room temperature. The mixture was filtered through a glass filter and the solvent was removed under reduced pressure. The crude product was column chromatographed (silica gel, 5→10% MeOH in CH<sub>2</sub>Cl<sub>2</sub>), to give 12.0 g 2-(N-CBz-L-valyloxy)-ethanol.

b) 2-(N-CBz-L-valyloxy)-ethyl chloromethyl carbonate

To a mixture of 2-(N-CBz-L-valyloxy)-ethanol (12.0 g, 40.6 mmol) and pyridine (19.7 mL, 0.24 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (300 mL) at 0 °C, was added dropwise chloromethyl chloroformate (10.5 g, 81.2 mmol). After 30 min at 0 °C, the reaction mixture was washed with H<sub>2</sub>O (200 mL). The H<sub>2</sub>O phase was washed with CH<sub>2</sub>Cl<sub>2</sub> (100 mL) and the solvent of the combined organic phases was removed under reduced pressure. The crude product was column chromatographed (silica gel, 0.5→1% MeOH in CH<sub>2</sub>Cl<sub>2</sub>), to give 8.26 g 2-(N-CBz-L-valyloxy)-ethyl chloromethyl carbonate.


c) 2-(N-CBz-L-valyloxy)-ethyl iodomethyl carbonate

To a solution of 2-(N-CBz-L-valyloxy)-ethyl chloromethyl carbonate (3.88 g, 10 mmol) in acetonitrile (50 mL), was added sodium iodide (7.50 g, 50 mmol). The solution was stirred for 4 h at 60 °C. The resulting suspension was filtered and the filtrate was evaporated. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and washed with aqueous sodium thiosulfate (0.1 M). The organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure, to give 4.51 g 2-(N-CBz-L-valyloxy)-ethyl iodomethyl carbonate.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 7.34 (s, 5H), 5.93 (s, 2H), 5.26 (d, 1H), 5.11 (s, 2H), 4.48-4.26 (m, 5H) 2.28-2.10 (m, 1H), 0.97 (d, 3H), 0.90 (d, 3H).

Example A-I-10

2,2-dimethyl-3-(N-CBz-D-valyloxy)-propionic acid iodomethyl ester



## a) 2,2-dimethyl-3-(N-CBz-D-valyloxy)-propionic acid

To a solution of 2,2-dimethyl propionic acid 4-methoxybenzyl ester (4.7 g, 20 mmole) and N-CBz-D-valine (5.5 g, 22 mmole) in dichloromethane (100 ml) were added 4-dimethylaminopyridine (305 mg, 2.5 mmole) and DCC (5.15 g, 25 mmole). After 18 hr, the solution was washed successively with sodium bicarbonate aqueous solution, citric acid solution and water. The organic phase was dried and the residue was dissolved in dichloromethane (100 ml). To the solution was added trifluoroacetic acid (10 ml). After 3 hr, it was evaporated and the product was isolated with silica gel column chromatography. 4.5 g.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 7.36 (m, 5 H) 5.11 (s, 2H) 4.30 (m, 1H) 4.18 (dd, 2H) 2.17 (m, 1H), 1.23 (d, 6 H) 0.93 (m, 6H).

b) 2,2-dimethyl-3-(N-CBz-D-Valyloxy)-propionic acid chloromethyl ester (2,2-dimethyl-3-(N-CBz-D-valyloxy)-propionic acid (4.5 g, 12.8 mmole) was dissolved in dioxane (20 ml). To the solution was added tetrabutylammonium hydroxide aqueous solution (40 %, 8.3 ml, 12.8 mmole). The solution was dried in vacuo, and it was coevaporated with toluene several times. The residue was dissolved in methylene chloride and then chloriodomethane (18 ml, 260 mmole) was added to the solution. After 18 hr, the reaction solution was evaporated and the product was isolated with silica gel column chromatography. 3.5 g.


c) 2,2-dimethyl-3-(N-CBz-D-valyloxy)-propionic acid iodomethyl ester 2,2-Dimethyl-3-(N-CBz-D-valyloxy)-propionic acid chloromethyl ester (2.4 g, 6 mmole) was dissolved in acetonitrile (30 ml). Sodium iodide (1.26 g, 8.4 mmole) was added to the solution. After reaction at 70° C for 2 hr, the reaction mixture was filtered and the residue was dissolved in methylene chloride (20 ml) and refiltered. The solution was dried and gave the titled product. 2.68g.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 7.36 (m, 5 H) 5.90 (dd, 2H) 5.26 (d, 1H) 5.11 (s, 2H) 4.31 (m, 1H) 4.15 (dd, 2H) 2.18 (m, 1H) 1.22 (d, 6H) 0.92 (m, 6H).

## Example A-1-11

4-(N-CBz-L-valyloxy) butyric acid iodomethyl ester





a) 4-(N-CBz-L-valyloxy) butyric acid t-butyl ester

N-CBz-L-valine (16.25 g, 65 mmole) was dissolved in DMF (40 ml). To the solution was added potassium t-butoxide (7.24 g, 65 mmole). After 10 min, 4-bromobutyric acid t-butyl ester (12 g, 53 mmole) was added. The reaction mixture was kept at 65° C for 2.5 hr and then poured into sodium bicarbonate aqueous solution and extracted with dichloromethane. The organic phase was dried and the product was isolated with silica gel column chromatography. 20.1 g.


b) 4-(N-CBz-L-valyloxy)butyric acid chloromethyl ester

4-(N-CBz-L-valyloxy) butyric acid t-butyl ester (20 g, 50.8 mmole) was treated with trifluoroacetic acid (30 ml) at 0° C for 3 h and then evaporated. The residue was coevaporated with toluene several time. The intermediate acid (2.56 g, 7.6 mmole) was dissolved in dioxane (10 ml) and to the solution was added tetrabutylammonium hydroxide (40 %, 4.66 ml, 7.2 mmole). The solution was dried and dissolved in dichloromethane (20 ml) and then chloriodomethane (10 ml, 144 mmole) was added to the solution. After 18 hr, the reaction solution was evaporated and the product was isolated with silica gel column chromatography. Yield 2.1 g.

c) 4-(N-CBz-L-valyloxy)butyric acid iodomethyl ester

4-(N-CBz-L-valyloxy) butyric acid chloromethyl ester (1.54 g, 4 mmole) was dissolved in acetonitrile (15 ml). Sodium iodide (840 mg, 5.6 mmole) was added to the solution. After reaction at 55° C for 3 hr, the reaction mixture was filtered and the residue was dissolved in methylene chloride (20 ml) and refiltered. The solution was dried and gave the titled product. Yield 1.9 g.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 7.36 (m, 5H) 5.90 (dd, 2 H) 5.25 (d, 1H) 5.11 (s, 2H) 4.29 (dd, 1H 4.18 (t, 2H) 2.43 (t, 2H) 2.20 (m, 1H) 2.00 (m, 2H) 0.93 (dd, 6 H).

Iodomethyl 3-(N-benzyloxycarbonyl-L-valyloxy)-benzoate

## a) 4-Methoxybenzyl 3-hydroxybenzoate

To a solution of 3-hydroxybenzoic acid (6.9g, 50 mmole) in DMF (100 ml)  
 5 was added potassium-tert.-butoxide (6.17 g, 55 mmole) and the mixture was  
 stirred at room temperature for one hour. 4-Methoxybenzyl chloride (9.4g, 60  
 mmole) was added and the mixture was stirred for 16 hours at 60°C. The  
 mixture was evaporated under reduced pressure and ethyl acetate (250 ml)  
 were added. The organic phase was washed five times with water, dried with  
 10 sodium sulfate and evaporated under reduced pressure. The product was  
 isolated by silica gel column chromatography with toluene/acetone. Yield:  
 10.5g = 81%

## b) 4-Methoxybenzyl 3-(N-benzyloxycarbonyl-L-valyloxy) benzoate .

To a cooled solution of 4-methoxybenzyl 3-hydroxybenzoate (7.7g, 29.8  
 15 mmole), 4-dimethylaminopyridine (0.73g, 6 mmole) and N-benzyloxycarbonyl-  
 L-valine (8.3g, 33 mmole) in 100 ml dichloromethane was added dicyclohexyl-  
 carbodiimide (7.22g, 35 mmole) and the mixture was stirred for 2 days at  
 room temperature. The mixture was cooled and the urethane was filtered. The  
 solution was evaporated and ethyl acetate (250 ml) was added. The organic  
 20 phase was washed twice with 5% acetic acid; 5% sodium hydrogencarbonate  
 and water. The organic phase was dried with sodium sulfate and evaporated  
 under reduced pressure. The product was isolated by silica gel column  
 chromatography with hexane/ethyl acetate. Yield: 13.9g = 94%

## c) 3-(N-benzyloxycarbonyl-L-valyloxy) benzoic acid

25 To a solution of 4-methoxybenzyl-3-(N-benzyloxycarbonyl-L-valyloxy)-  
 benzoate (13.7g, 27.8 mmole) in dichloromethane (150 ml) was added  
 trifluoroacetic acid (20 ml) and the mixture was stirred for 2 hours at room  
 temperature. The solution was evaporated under reduced pressure and the  
 product crystallized from toluene. Yield: 10.1g = 87%.  
 30 <sup>1</sup>H-NMR (DMSO d-6) 1.01 (m, 6H) 2.21 (m, 1H) 4.17 (d, d, 1H) 5.08 (s, 2H)  
 7.28-7.96 (m, 10H)

d) Chloromethyl 3-(N-benzyloxycarbonyl-L-valyloxy)-benzoate.

To a solution of 3-(N-benzyloxycarbonyl-valyloxy)benzoic acid (7.42g, 20 mmole) in 1,4-dioxane (100 ml) was added a 40% solution of tetrabutylammonium hydroxide (12.97g, 20 mmole) and the mixture was stirred 2 hours at room temperature. The mixture was evaporated under reduced pressure and co-evaporated two times with 1,4-dioxane and two times with toluene. The dried product was dissolved in dichloromethane (50 ml) and chloriodomethane (35.3g, 200 mmole) was added. The solution was stirred for two days at room temperature and evaporated under reduced pressure. Ethyl acetate (100 ml) was added and the organic phase washed twice with water, dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography. Yield: 3.8g = 45%.


e) Iodomethyl 3-(N-benzyloxycarbonyl-L-valyloxy)-benzoate.

To a solution of chloromethyl 3-(N-benzyloxycarbonyl-L-valyloxy)-benzoate (2.0g, 4.76 mmole) in dry acetone (30 ml) was added sodium iodide (3.15g, 21 mmole) and the mixture was stirred overnight at room temperature. The mixture was evaporated under reduced pressure and extracted with ethyl acetate/water. The organic phase was washed with a 5% sodium thiosulfate solution, dried with sodium sulfate and evaporated under reduced pressure. Yield: 2.3g = 94%.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 1.02 (m, 6H) 2.38 (m, 1H) 4.56 (d, d, 1H) 5.14 (s, 2H) 5.30 (d, 1H) 6.14 (s, 2H) 7.26-7.50 (m, 7H) 7.80(s, 1H) 7.96 (d, 1H)

#### Example A-I-13

##### Iodomethyl 3-(N-benzyloxycarbonyl-L-valyloxy)-propionate



a) 3-buten-1-yl-3-(N-benzyloxycarbonyl)-propionate.

To a solution of 3-buten-1-ol (2.16g, 30 mmole), N-benzyloxycarbonyl-L-valine (8.29g, 33 mmole) and 4-dimethylaminopyridine (0.37g, 3 mmole) in

dichloromethane (80 ml) was added dicyclohexyl-carbodiimide (7.22g, 35 mmole) and the mixture was stirred overnight at room temperature. The mixture was cooled and the urethane was filtered. The solution was evaporated under reduced

- 5 pressure and ethyl acetate (200 ml) was added. The organic phase was washed twice with 5% acetic acid, 5% sodium hydrogencarbonate and water. The organic phase was dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography with hexane/ethyl acetate. Yield: 8.3g = 90%.

10 b) 3-(N-benzyloxycarbonyl-L-valyloxy)-propanoic acid

To a solution of 3-buten-1-yl -3-( N-benzyloxycarbonyl-L-valyloxy)-propionate (9.2g, 30 mmole) in 150 ml benzene was added tetrabutylammonium bromide (1.62g, 5 mmole) and 100 ml water. The mixture was cooled to about 5°C and potassium permanganate (14.82g, 90 mmole) was added in portions. The

- 15 mixture was stirred 2 hours at room temperature, diluted with water and decolorized by the addition of sodium bisulfite. The mixture was acidified with 2M hydrogen chloride and extracted 3 times with ethyl acetate. The combined organic phases were washed with water and dried with sodium sulfate. The solution was evaporated under reduced pressure and the product isolated by  
20 silica gel column chromatography with hexane/ethyl acetate. Yield: 5.4g = 55%.

<sup>1</sup>H-NMR (DMSO d-6) 0.90 (m, 6H) 2.5 (m, 2H) 3.88 (d, d, 1H) 4.32 (m, 2H) 5.03 (s, 2H) 7.36 (m, 5H) 7.68 (d, 1H)

c) Chloromethyl 3-(N-benzyloxycarbonyl-L-valyloxy)-propionate.

- 25 To a solution of 3-(N-benzyloxycarbonyl-L-valyloxy)propanoic acid (5.2g, 16.08 mmole) in 1,4-dioxane (50 ml) was added a 40% solution of tetrabutylammonium hydroxide (10.43g, 16.08 mmole) and the mixture was stirred 2 hours at room temperature. The mixture was evaporated under reduced pressure and co-evaporated two times with 1,4-dioxane and two  
30 times with toluene. The dried product was dissolved in 40 ml dichloromethane and chloriodomethane (28.4g, 160 mmole) was added. The solution was stirred for two days at room temperature and evaporated under reduced pressure. Ethyl acetate (100 ml) was added and the organic phase washed twice with water, dried with sodium sulfate and evaporated under reduced

pressure. The product was isolated by silica gel column chromatography.

Yield: 2.2g = 35%

d) Iodomethyl 3-(N-benzyloxycarbonyl-L-valyloxy)-propionate.


To a solution of chloromethyl 3-(N-benzyloxycarbonyl-L-valyloxy)-propionate  
5 (2.05g, 5.51 mmole) in dry acetone (50 ml) was added sodium iodide (4.12g,  
27.5 mmole) and the mixture was stirred overnight at room temperature. The  
mixture was evaporated under reduced pressure and extracted with ethyl  
acetate water. The organic phase was washed with a 5% sodium thiosulfate  
solution, dried with sodium sulfate and evaporated under reduced pressure.

10 Yield: 2.35g = 92%.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.94 (m, 6H) 2.17 (m, 1H) 2.68 (t, 2H) 4.40 (m, 3H) 5.12 (s,  
2H) 5.91 (s, 2H) 7.26 (m, 5H).

Example A-I-15

15 1-(N-benzyloxycarbonyl-L-valyloxy)-2-methyl-2-propyl iodomethyl carbonate



(a) 1-(N-benzyloxycarbonyl-L-valyloxy)-2-methyl-2-propanol

N-Benzyloxycarbonyl-L-valine (2.02 g, 8.0 mmol), 4-dimethylaminopyridine  
(100 mg, 0.8 mmol), and ), and dicyclohexylcarbodiimide (2.04 g, 9.9 mmol, in  
20 20 mL CH<sub>2</sub>Cl<sub>2</sub>) were added to 2-methyl-1,2-propanediol (12.2 mmol) in 30 mL  
dry CH<sub>2</sub>Cl<sub>2</sub>, with cooling in an ice bath. DMF (5 mL) was added. After stirring  
for 5 h at 10 °C , the reaction mixture was filtered, concentrated, and then  
redissolved in ethyl acetate. The organic solution was washed with saturated  
NaCl, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, and concentrated. Flash column  
25 chromatography (silica, 2/1 petroleum ether – ethyl acetate) gave 2.3 g of the  
title compound.

(b) 1-(N-benzyloxycarbonyl-L-valyloxy)-2-methyl-2-propyl chloromethyl  
carbonate

All of the alcohol from above was dissolved in 35 mL dry CH<sub>2</sub>Cl<sub>2</sub> and cooled in  
30 an ice bath. Pyridine (3.50 mL, 43.4 mmol) was added, followed by

chloromethyl chloroformate (1.30 mL, 14.4 mmol). After 1 h, the ice bath was removed and stirring was continued for 2 h at ambient temperature. The mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (50 mL) and washed with water (50 mL), and then brine (2 x 25 mL). Drying over anhydrous  $\text{Na}_2\text{SO}_4$  of the combined  
 5 organic phases and concentration under vacuum, coevaporating several times with toluene, gave a yellow-brown oil that was subjected to flash column chromatography (silica, 2/1 petroleum ether – ethyl acetate) to yield 2.86 g (86% from *N*-benzyloxycarbonyl-L-valine) of the title compound.


c) 1-(*N*-benzyloxycarbonyl-L-valyloxy)-2-methyl-2-propyl iodomethyl  
 10 carbonate

A mixture of the chloride (2.84 g, 6.84 mmol) from step (b) and NaI (4.15 g, 27.2 mmol) in 68 mL dry acetonitrile was refluxed at 75 °C for 4 h. After evaporation of solvent under vacuum, the residue was partitioned between ethyl acetate (80 mL) and water (40 mL), and the organic layer was washed  
 15 with 5%  $\text{Na}_2\text{S}_2\text{O}_3$  (15 mL) and brine (25 mL). Drying the organic phase over anhydrous  $\text{Na}_2\text{SO}_4$  and concentration gave a yellow oil that was subjected to flash column chromatography (silica, 2/1 petroleum ether - ethyl acetate) to furnish 3.29 g (95%) of the title compound.

20  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ ) 0.90 and 0.94 (2d, 3H each,  $J = 6.8$  Hz), 1.52 (s, 6H), 2.17 (m, 1H), 4.35 (m, 1H), 4.22 and 4.39 (ABq, 2H,  $J_{AB} = 11.7$  Hz), 5.10 (s, 2H), 5.30 (d, 1H), 5.86 (s, 2H), 7.34 (s, 5H)

#### Example A-I-16

25 Iodomethyl 3,4-di-(*N*-CBZ-L-valyloxy)hydrocinnamate



a) 4-Methoxybenzyl-3,4-dihydroxyhydrocinnamate

3,4-Dihydroxycinnamic acid (6.5 g, 35.7 mmol) was dissolved in DMF (50 ml) and cooled to 0°C on an ice-bath. 4-Potassium tert-butoxide (35.7 mmol), was

then added and the mixture was left for approximately 30 min at 0°C, followed by dropwise addition of 4-methoxy-benzylchloride (39 mmol) in DMF (25 ml).

The mixture was allowed to reach room temperature and left over-night. The solvent was then evaporated and the crude product was purified by

5 chromatography (ethyl acetate-hexane, 1:1) to give 6 g of the title compound (55%).

b) 4-Methoxybenzyl-3,4-di-(N-CBZ-L-valyloxy)hydrocinnamate

4-Methoxybenzyl-3,4-dihydroxyhydrocinnamate (5 g, 16.5 mmol), *N,N*-dimethylaminopyridine (2g, 16.5 mmol), *N,N'*-dicyclohexyl carbodiimide (8.5

10 g, 41.3 mmol) and Cbz-L-valine (10.4 g, 41.3 mmol) were dissolved in

dichloromethane (50 ml). After 4 h, the mixture was filtered and evaporated onto silica gel and purified by chromatography (hexane-EtOAc,

5:2 → 3:2) to give pure title product (10.1 g, 79 %).

c) 3,4-Di-(N-CBZ-L-valyloxy)hydrocinnamic acid

15 4-Methoxybenzyl-3,4-di-(N-CBZ-L-valyloxy)hydrocinnamate (10 g, 13 mmol)

was dissolved in dichloromethane and 1,1,1 trifluoroacetic acid (30 ml) and

left at ambient temperature for 3.5 h. Evaporation under reduced pressure and

purification by chromatography (chloroform-methanol, 10:1) yielded 6.7 g

(80%) pure title product.

20 <sup>1</sup>H NMR (CDCl<sub>3</sub> 45 °C): 7.24-7.0 (m, 13H), 5.65 (br s, 1H), 5.55 (br s, 1H), 5.1 (m, 4H), 4.46 (m, 2H), 2.95 (t, 2H), 2.66 (t, 2H), 2.35 (m, 2H).

d) Chloromethyl 3,4-di-(N-CBZ-L-valyloxy)hydrocinnamate

3,4-Di-(N-CBZ-L-valyloxy)hydrocinnamic acid (4.2 g, 6.47 mmol) was

dissolved in dioxane (70 ml). Tetrabutylammonium hydroxide was added

25 dropwise until pH=8. The solvent was then removed under reduced pressure

The solid was redissolved in dioxane (30 ml) and toluene (30 ml) and

evaporated. The procedure was repeated twice (for removal of water).

Dichloromethane (60 ml) and chloro-iodomethane was added in one portion

and the mixture was left at ambient temperature for 6 h. Evaporation of the

30 solvent and purification by chromatography yielded 1.7 g title product (38%).

e) Iodomethyl 3,4-di-(N-CBZ-L-valyloxy)hydrocinnamate

Chloromethyl 3,4-di-(N-CBZ-L-valyloxy)hydrocinnamate (1.9 g, 2.7 mmol) and


sodium iodide (2 g, 13.3 mmol) were dissolved in acetonitrile (50 ml) and

heated to 65° C for 60 min. The solvent was removed under reduced pressure and the residue was taken up in dichloromethane and filtrated. Removal of the solvent and purification by chromatography (ethyl acetate-hexane, 2:5) gave pure title product (1.9 g, 90 %).

5

<sup>1</sup>H NMR (CDCl<sub>3</sub> 45 °C): 7.34-7.02 (m, 13H), 5.89 (s, 2H), 5.64 (br s, 2H), 5.14-5.02 (m, 4H), 4.47 (m, 2H), 2.96 (t, 2H), 2.64 (t, 2H), 2.33 (m, 2H), 1.08-0.99 (m, 12H)

## 10 Example A-I-17

3-(N-CBz-L-valyloxy)phenyl iodomethyl carbonate

## a) 3-(N-CBz-L-valyloxy)phenol.

CBz-L-valine (10 g, 40 mmol), 1,3-dihydroxybenzene (8.7g, 79 mmol)

15 *N,N*'dicyclohexylcarbodiimide (10.2g, 44 mmol) and 4-dimethylaminopyridine (2.4 g, 20 mmol) were dissolved in DMF (50 ml) and left at ambient temperature overnight. The reaction mixture was filtered, the solvent removed under reduced pressure and the crude product was taken up in dichloromethane and filtered. Removal of the solvent followed by purification  
20 by chromatography (chloroform-methanol, 10:1) yielded pure title product (10.9 g, 79%).

## b) (N-CBz-L-valyloxy)phenyl chloromethyl carbonate.

3-(N-CBz-L-valyloxy)phenol (5.4 g, 15.7 mmol) was dissolved in dichloromethane (70 ml) and cooled in an ice-bath. Pyridine (1.2 g, 23.5 mmol)  
25 was added followed by dropwise addition of 1-chloro-methylchloroformate (2.3 g, 18.8 mmol) in dichloromethane (10 ml). The mixture was left at room temperature for 4 h. Water (25 ml) was then added and the phases were separated. The organic layer was washed with 0.01 M aqueous hydrochloric acid (25 ml). Purification by chromatography (ethyl acetate-hexane, 1:1) gave  
30 the title compound (4.5 g, 65 %)




c) 3-(N-CBZ-L-valyloxy)phenyl iodomethyl carbonate

(N-CBZ-L-valyloxy)phenyl chloromethyl carbonate (1.5g, 3.44 mmol) and sodium iodide (2 g, 13.3 mmol) were stirred at 60°C in acetonitrile (50 ml) for 4.5 h. The mixture was filtered, the solvent removed and the crude product was taken up in 100 ml hexane-ethyl acetate, 1:1, and filtered through a sintered glass funnel, packed with 2 cm silica gel. Removal of the solvent yielded pure title product (1.68 g, 92%)

$^1\text{H}$  NMR ( $\text{CDCl}_3$ , 45 °C): 7.38-7.02 (m, 9H), 6.03 (s, 2H), 5.2 (br s, 1H), 5.14 (s, 2H), 4.48 (m, 1H), 2.30 (m, 1H), 1.09-1.01 (m, 6H)

#### Example A-I-18

##### Iodomethyl 2-(N-CBZ-L-valyloxy)phenylacetate



a) 4-Methoxybenzyl 2-hydroxyphenylacetate.

2-hydroxyphenylacetic acid (10 g, 66 mmol) was dissolved in *N,N*-dimethyl-formamide (100 ml) and cooled on ice-bath. Potassium *tert*-butoxide (8.85 g, 78 mmol) was added. The mixture was left for 30 min and allowed to reach room temperature. 4-Methoxy-benzylchloride (11.7 g, 72 mmol) in *N,N*-dimethyl-formamide (30 ml) was then added dropwise, under nitrogen atmosphere and left over-night. The solvent was evaporated under reduced pressure and the crude mixture was dissolved in ether (100 ml) and washed with water (25 ml), brine and dried over sodium sulphate. Chromatography (hexane-ethyl acetate, 2:1) followed by recrystallization (hexane-ethyl acetate) gave the title compound (7.6 g, 42%).

b) 4-Methoxybenzyl 2-(N-CBZ-L-valyloxy)phenylacetate

4-Methoxybenzyl 2-hydroxyphenylacetate 3g, 11 mmol), *N,N*-dichyclohexyl-carbodiimide (2.7 g, 13.2 mmol), dimethylaminopyridine (0.134 g, 1.1 mmol) and CBZ-L-valine (3.3 g, 13.2 mmol) were dissolved in dichloromethane (50 ml). After the weekend the solid was filtered off, the solvent removed under

reduced pressure and the crude product purified by chromatography (ethyl acetate, hexane, 1:2) to give the title compound (5.2 g, 93%).

c) 2-(N-CBz-L-valyloxy)phenylacetic acid

4-Methoxybenzyl 2-(N-CBz-L-valyloxy)phenylacetate (4.25 g, 8.4 mmol), was dissolved in dichloromethane (40 ml). Trifluoroacetic acid (8 ml) was added with cooling on ice. The mixture was allowed to reach room temperature and stirred for 40 min. The solvent was removed under reduced pressure and the crude product was recrystallized twice (hexan-ethyl acetate + a small amount of dichloromethane) to give the title compound (2.6 g, 80 %).

<sup>1</sup>H NMR (CDCl<sub>3</sub> 45 °C): 7.35-7.08 (m, 9H), 5.35 (br s, 1H), 5.13 (s, 2H), 4.48 (m, 1H), 3.57 (s, 2H), 2.33 (m, 1H), 1.08 (d, 3H), 1.02 (d, 3H).

d) Chloromethyl 2-(N-CBZ-L-valyloxy)phenylacetate

This compound was prepared in poor yield from 2-(N-CBZ-L-valyloxy)phenylacetic acid (5.5 g, 14.3 mmol) by an unoptimized procedure essentially as described in Example A-I-16 d). Yield: 0.265 g


<sup>1</sup>H NMR (CDCl<sub>3</sub> 45 °C): 7.28-7.01 (m, 9H), 5.55 (s, 2H), 5.2 (br s, 1H), 5.07 (s, 2H), 4.43 (m, 1H), 3.53 (s, 2H), 2.26 (m, 1H), 1.02 (d, 3H), 0.95 (d, 3H).

e) Iodomethyl 2-(N-CBZ-L-valyloxy)phenylacetate

Chloromethyl 2-(N-CBZ-L-valyloxy)phenylacetate is treated with NaI and purified as described in the Examples above to yield the title compound.

Example A-I-19

Iodomethyl 4-(N-CBZ-L-valyloxyxy)phenylacetate



25

a) 4-Methoxybenzyl 4-hydroxyphenylacetate

Prepared from 4-hydroxyphenylacetic acid (10 g, 65.7 mmol) in 70 % yield by the same procedure as for Example A-I-18 a) above, but wherein the solvent for the recrystallization was changed to hexane-ether.

## b) 4-Methoxybenzyl 4-(N-CBz-L-valyloxy)phenylacetate

Prepared from 4-methoxybenzyl 4-hydroxyphenylacetate (3 g, 11 mmol) by the same procedure as for Example A-I-18 b) in 87 % yield. Solvent for chromatography: ethyl acetate-hexane, 1:2.

## 5 c) 4-(N-CBZ-L-valyloxy)phenylacetic acid

Prepared in 82 % yield from 4-methoxybenzyl 4-(N-CBz-L-valyloxy)phenylacetate (1.6 g, 288 mmol) by the procedure described for Example A-I-18 c). Solvent for recrystallization: hexane-ether and a small amount of dichloromethane.

10  $^1\text{H}$  NMR ( $\text{CDCl}_3$  45 °C): 7.36-7.27 (m, 7H), 7.02 (d, 2H), 5.25 (d, 1H), 5.14 (s, 2H), 4.52 (m, 1H), 3.64 (s, 2H), 2.3 (m, 1H), 1.08 (d, 3H), 1.02 (d, 3H).

## d) Chloromethyl 4-(N-CBZ-L-valyloxy)phenylacetate

Prepared from 4-(N-CBZ-L-valyloxy)phenylacetic acid (3 g, 7.8 mmol) in 26 % yield by the same procedure as described for Example A-I-18 d). Solvent for  
15 chromatography: hexane-ether, 3:2.

## e) Iodomethyl 4-(N-CBZ-L-valyloxy)phenylacetate

Chloromethyl 4-(N-CBZ-L-valyloxy)phenylacetate (0.83 g, 1.9 mmol) and sodium iodide (1.15 g, 7.6 mmol) were heated in acetonitril (45 ml) for 5 h. The mixture was filtrated, the solvent removed, taken up in dichloromethane  
20 and filtrated again. Evaporation and purification by chromatography (ether-hexane, 2:3) yielded the title product (0.8 g, 80 %).

$^1\text{H}$  NMR ( $\text{CDCl}_3$  45 °C): 7.38-7.09 (m, 4H), 5.84 (s, 1H), 5.30 (br s, 1H), 5.15 (s, 2H), 4.5 (m, 1H), 3.56 (s, 2H), 2.36 (m, 1H), 1.10 (d, 3H), 1.00 (d, 3H).

25

## Example A-I-20

Iodomethyl 4-(2-N-benzyloxycarbonyl-L-valyloxyethyl) benzoate

## a) 4-(2-N-benzyloxycarbonyl-L-valyloxyethyl)-toluene

30 To a cooled solution of 4-methylphenylethanol-2 (5.0g, 36.7 mmole), 4-dimethylaminopyridine (0.98g, 8 mmole) and N-benzyloxycarbonyl-L-valine (10.05g, 40 mmole) in dichloromethane (120 ml) was added dicyclohexylcarbodiimide (9.1g, 44 mmole) and the mixture was stirred overnight at room temperature. The mixture was cooled and the urethane was filtered. The

solution was evaporated under reduced pressure and ethyl acetate (250 ml) was added. The organic phase was washed twice with 5% acetic acid, 5% sodium hydrogencarbonate and water. The organic phase was dried with sodium sulfate and evaporated under reduced pressure. The product was  
5 isolated by silica gel column chromatography

b) 4-(2-N-benzyloxycarbonyl-L-valyloxyethyl)- benzoic acid .

To a cooled mixture of chromic anhydride (7.55g, 75 mmole) in acetic acid (100 ml) was added dropwise a solution of 4-(2-N-benzyloxycarbonyl-L-valyloxyethyl)-toluene (9.3g, 25.1 mmole) in acetone (50 ml). The mixture was  
10 stirred at room temperature for 3 days and reduced to about 100 ml. 600ml 10% sodium chloride solution was added and the mixture was extracted four times with ethyl acetate. The organic phase was washed with brine and dried with sodium sulfate. The solution was evaporated under reduced pressure and the product was isolated by silica gel column chromatography with  
15 dichloromethane/methanol. Yield : 2.1g = 21%.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.79 (d, 3H) 0.90 (d, 3H) 2.08 (m, 1H) 3.04 (t, 2H) 4.28 (d, 1H) 4.39 (m, 2H) 5.11 (s, 2H) 5.26 (d, 1H) 7.34 (m, 7H) 8.04 (d, 2H)

c) Chloromethyl 4-(2-N-benzyloxycarbonyl-L-valyloxyethyl)benzoate

To a solution of 4-(2-N-benzyloxycarbonyl-L-valyloxyethyl)benzoic acid (2.0g, 5.0 mmole) in 1,4-dioxane (20 ml) was added a 40% solution of  
20 tetrabutylammonium hydroxide (3.1g, 4.75 mmole) and the mixture was stirred 2 hours at room temperature. The mixture was evaporated under reduced pressure and coevaporated two times with 1,4-dioxane and two times with toluene. The dried product was dissolved in dichloromethane (10 ml) and  
25 iodochloromethane (13.2g, 75 mmole) was added. The solution was stirred overnight at room temperature and evaporated under reduced pressure. About 50 ml ethyl acetate were added and the organic phase washed twice with water, dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography. Yield: 0.5g  
30 =23%

d) Iodomethyl 4-(2-N-benzyloxycarbonyl-L-valyloxyethyl) benzoate

To a solution of chloromethyl 4-(2-N-benzyloxycarbonyl-L-valyloxyethyl) benzoate (0.5g, 1.11 mmole). In dry acetone (10 ml) was added sodium iodide (0.75g, 5.0 mmole) and the mixture was stirred overnight at room

temperature. The mixture was evaporated under reduced pressure and extracted with ethyl acetate/water. The organic phase was washed with a 5% sodium thiosulfate solution, dried with sodium sulfate and evaporated under reduced pressure. Yield: 0.53g = 88%.

5

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.88 (d, 3H) 0.90 (d, 3H) 2.08 (m, 1H) 3.02 (t, 2H) 4.28 (d, d, 1H) 4.38 (m, 2H) 5.10 (s, 2H) 5.22 (d, 1H) 6.15 (s, 2H) 7.35(m, 7H) 7.98 (d, 2H )

## 10 Example A-I-21

Iodomethyl 2-(N-benzyloxycarbonyl-L-isoleucyloxymethyl)-  
2-methyl propionate.

a) 4-methoxybenzyl 2-(N-benzyloxycarbonyl-L-isoleucyloxymethyl)-  
2-methyl propionate

15 To a cooled solution of 4-methoxybenzyl 2-(hydroxymethyl)-2-methyl  
propionate (6.0g, 25 mmole). 4-dimethylaminopyridine (0.61g, 5 mmole) and  
N-benzyloxycarbonyl-L-isoleucine (6.90g, 26 mmole) in dichloromethane (100  
ml) was added dicyclohexyl-carbodiimide (6.2g, 30 mmole) and the mixture  
was stirred overnight at room temperature. The mixture was cooled and the  
20 urethane was filtered. The solution was evaporated and 200 ml ethyl acetate  
was added, The organic phase was washed twice with 5% acetic acid, 5%  
sodium hydrogencarbonate and water. The organic phase was dried with  
sodium sulfate and evaporated under reduced pressure. The product was  
isolated by silica gel column chromatography with toluene/acetone. Yield:

25 11.7g = 96%.

2-(N-benzyloxycarbonyl-L-isoleucyloxymethyl)-2-methyl) propionic acid.

To a solution of 4-methoxybenzyl 2-(N-benzyloxycarbonyl-L-  
isoleucyloxymethyl)-2-methyl propionate (11.0g, 22.6 mmole) in 100 ml  
dichloromethane was added trifluoroacetic acid (15 ml) and the mixture was  
30 stirred overnight at room temperature. The solution was evaporated under  
reduced pressure and coevaporated two times with toluene. The residue was  
stirred 1 hour with 100 ml ethanol and the white solid was filtered (byproduct).  
The solution was evaporated under reduced pressure and the product was

isolated by silica gel column chromatography with hexane/ethyl acetate. Yield: 7.4g = 89%.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.90 (m, 6H) 1.26 (m, 8H) 1.88 (m, 1H) 4.12 (d, d, 2H) 4.38 (d, d, 1H) 5.10 (s, 2H) 5.32 (d, 1H) 7.28 (m, 5H)

- 5 c) Chloromethyl 2-(N-benzyloxycarbonyl-L-isoleucyloxy)-2-methyl propionate.

To a solution of 2-(N-benzyloxycarbonyl-L-isoleucyloxymethyl)-2-methyl propionic acid (7.0g, 19 mmole) in 80 ml 1,4-dioxane was added a 40% solution of tetrabutylammonium hydroxide (12.4g, 19 mmole) and the mixture  
10 was stirred 2 hours at room temperature. The mixture was evaporated under reduced pressure and co-evaporated two times with 1,4-dioxane and two times with toluene. The dried product was dissolved in 25 ml dichloromethane and iodochloromethane (33.7g, 190 mmole) was added. The solution was stirred overnight at room temperature and evaporated under reduced  
15 pressure. About 100 ml ethyl acetate was added and the organic phase washed twice with water, dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography with toluene/acetone. Yield: 4.2 = 54%

- 20 d) Iodomethyl 2-(N-benzyloxycarbonyl-L-isoleucyloxymethyl)-2-methyl propionate.

To a solution of chloromethyl 2-(N-benzyloxycarbonyl-L-isoleucyloxymethyl)-2-methyl propionate (3.0g, 7.2 mmole) in 50 ml dry acetone was added sodium iodide (4.8g, 32 mmole) and the mixture was stirred overnight at room temperature. The mixture was evaporated under reduced pressure and  
25 extracted with ethyl acetate water. The organic phase was washed with a 5% sodium thiosulfate solution, dried with sodium sulfate and evaporated under reduced pressure. Yield: 3.3g = 90%.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.93 (m, 6H) 1.23 (m, 8H) 4.12 (m, 2H) 4.38 (d, d, 1H) 5.10  
30 (s, 2H) 5.26 (d, 1H) 5.92 (m, 2H) 5.35 (m, 5H)

#### Example A-I-22

Iodomethyl 4-(N-benzyloxycarbonyl-L-valyloxy)cyclohexanoate.

a) 4-Methoxybenzyl 4-hydroxycyclohexanoate.

To a solution of ethyl 4-hydroxycyclohexanoate (8.61g, 50 mmole) in 50 ml ethanol was added a solution of potassium hydroxide 85% (3.63g, 55 mmole) and the mixture was stirred for 6 hours at 70°C. The mixture was evaporated under reduced pressure, coevaporated two times with N,N-dimethylformamide and reduced to about 100 ml. 4-Methoxybenzyl chloride (9.4g, 60 mmole) was added and the mixture was stirred for 18 hours at 60°C. The mixture was evaporated under reduced pressure and 250 ml ethyl acetate was added. The organic phase was washed five times with water, dried with sodium sulfate and evaporated under reduced pressure. Yield: 13.2g =100% (crude)

b) 4-methoxybenzyl 4-(N-benzyloxycarbonyl-L-valyloxy)-cyclohexanoate.

To a cooled solution of 4-methoxybenzyl 4-hydroxycyclohexanoate (7.5g, 28 mmole), 4-dimethylaminopyridine (0.73g, 6 mmole) and N-benzyloxycarbonyl-L-valine (7.54g, 30 mmole) in dichloromethane (90 ml) was added dicyclohexylcarbodiimide (6.8g, 33 mmole) and the mixture was stirred for 2 days at room temperature. The mixture was cooled and the urethane was filtered. The solution was evaporated and 250 ml ethyl acetate was added. The organic phase was washed twice with 5% acetic acid, 5% sodium hydrogencarbonate and water. The organic phase was dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography with toluene/acetone. Yield : 13g = 93%

c) 4-(N-benzyloxycarbonyl-L-valyloxy) cyclohexanoic acid.

To a solution of 4-methoxybenzyl 4-(N-benzyloxycarbonyl-L-valyloxy)-cyclohexanoate (12g, 24.1 mmole) in dichloromethane (100 ml) was added trifluoroacetic acid (20 ml) and the mixture was stirred for 3 hours at room temperature. The solution was evaporated under reduced pressure and coevaporated two times with toluene. The residue was stirred 1 hour with about 100 ml ethanol and the white solid was filtered (byproduct). The solution was evaporated under reduced pressure and the product was isolated by silica gel column chromatography with toluene/acetone. Yield: 6.8g = 74%.  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.91 (m, 6H) 1.52-2.54 (m, 10H) 4.28 (m, 1H) 4.82-5.08 (m, 1H) 5.11 (s, 2H) 5.28 (d, 1H) 7.36 (m, 5H)

d) Chloromethyl 4-(N-benzyloxycarbonyl-L-valyloxy)-cyclohexanoate.

To a solution of 4-(N-benzyloxycarbonyl-L-valyloxy) cyclohexanoic acid (6.6g, 20 mmole) in 1,4-dioxane (70 ml) was added a 40% solution of tetrabutylammonium hydroxide (11.34g, 17.5 mmole) and the mixture was stirred 2 hours at room temperature. The mixture was evaporated under reduced pressure and co-evaporated two times with 1,4-dioxane and two times with toluene. The dried product was dissolved in 60 ml dichloromethane and iodochloromethane (30.9g, 175 mmole) was added. The solution was stirred for two days at room temperature and evaporated under reduced pressure. About 100 ml ethyl acetate was added and the organic phase washed twice with water, dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography with toluene/acetone. Yield: 4.1g = 55%.


e) Iodomethyl 4-(N-benzyloxycarbonyl-L-valyloxy)-cyclohexanoate.

To a solution of chloromethyl 4-(N-benzyloxycarbonyl-L-valyloxy)-cyclohexanoate (4.0g, 9.4 mmole) in dry acetone (50 ml) was added sodium iodide (6.3g, 42 mmole) and the mixture was stirred overnight at room temperature. The mixture was evaporated under reduced pressure and extracted with ethyl acetate water. The organic phase was washed with a 5% sodium thiosulfate solution, dried with sodium sulfate and evaporated under reduced pressure. Yield 4.5g = 93%.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.90 (m, 6H) 1.52-2.02 (m, 8H) 2.18 (m, 1H) 2.43 (m, 1H) 4.30 (m, 1H) 4.76-5.08 (m, 1H) 5.11 (s, 2H) 5.26 (d, 1H) 5.91 (d, 2H) 7.34 (m, 5H)

#### Example A-I-23

#### Iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-ethyl butyrate





## a) 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-ethylbutan-1-ol.

To a cooled solution of 2-ethyl-2-hydroxymethyl-butan-1-ol (33.1g, 250 mmole), 4-dimethylaminopyridine (1.22g, 10 mmole) and N-benzyloxycarbonyl-L-valine (12.6g, 50 mmole) in 350 ml dichloromethane was added dropwise a solution of dicyclohexyl-carbodiimide (12.4g, 60 mmole) in 50 ml dichloromethane. The mixture was stirred 2 days at room temperature and cooled. The urethane was filtered and the solution evaporated under reduced pressure. 350 ml ethyl acetate was added and the organic phase was washed twice with 5% acetic acid, 5% sodium-hydrogencarbonate and water. The organic phase was dried with sodium sulfat and evaporated under reduced pressure. The product was isolated by silica gel column chromatography with dichloromethane/methanol. Yield: 16.4g = 90%.

## c) 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-ethyl-butyric acid.

To a cooled mixture of chromic anhydride (8.5g, 85,2 mmole) in 100 ml acetic acid was added dropwise a solution of 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-ethyl-butan-1-ol (10.4g, 28.4 mmole) in 50 ml acetone and the mixture was stirred 24 hours at room temperature. The mixture was added to 1000 ml 10% sodium chloride solution and extracted four times with ethyl acetate. The organic phase was washed twice with brine, dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography with hexane/ethyl acetate. Yield: 7g = 65%. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.88 (m, 12H) 1.67 (m, 4H) 2.14 (m, 1H) 4.26 (m, 3H) 5.10 (s, 2H) 5.30 (d, 2H) 7.34 (m, 5H)

## d) Chloromethyl 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-ethyl butyrate.

To a solution of 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-ethyl-butyric acid (7.2g, 18,9 mmole) in 1,4-dioxane (80 ml) was added a 40% solution of tetrabutylammonium hydroxide (12.26g, 18.9 mmole) and the mixture was stirred 2 hours at room temperature. The mixture was evaporated under reduced pressure and co-evaporated once with 1,4-dioxane and two times with toluene. The dried product was dissolved in 30 ml dichloromethane and iodochloromethane (49.4g, 280 mmole) was added. The solution was stirred for two days at room temperature and evaporated under reduced pressure. About 100 ml ethyl actate were added and the organic phase washed twice

with water, dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography. Yield: 5.2g = 63%

e) Iodomethyl 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-ethyl butyrate.


- 5 To a solution of chloromethyl 2-(N-benzyloxycarbonyl-L-valyloxymethyl)-2-ethyl butyrate (5.0g, 11.7 mmole) in dry acetone (60 ml) was added sodium iodide (7.5g, 50 mmole) and the mixture was stirred overnight at room temperature. The mixture was evaporated under reduced pressure and extracted with ethyl acetate water. The organic phase was washed with a 5% sodium thiosulfate solution, dried with sodium sulfate and evaporated under reduced pressure. Yield: 5.4g = 90%.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.92 (m, 12H) 1.65 (m, 4H) 2.18 (m, 1H) 4.28 (m, 3H) 5.10 (s, 2H) 5.22 (d, 1H) 5.92 (s, 2H) 7.36 (m, 5H)

15

Example A-I-24

2-(N-(iodomethoxycarbonyl)-amino)-2-methyl-1-(N-benzyloxycarbonyl-L-valyloxy)-propane



20

a) 2-(N-tert.-butyloxycarbonylamino)-2-methyl-1-(N-benzyloxycarbonyl-L-valyloxy)-propane.

- To a cooled solution of 2-(N-(tert.-butyloxycarbonyl)-amino)-2-methylpropan-1-ol (J. Am. Chem. Soc 113 (1991) p 8883) (4.73g, 25 mmole), 4-dimethylamino-pyridine (0.61g, 5 mmole) and N-benzyloxycarbonyl-L-valine (6.28g, 25 mmole) in dichloromethane (70 ml) was added dicyclohexyl-carbodiimide (6.19g, 30 mmole) and the mixture was stirred 2 days at room temperature. The mixture was cooled, the urethane was filtered and the solution evaporated under reduced pressure. Ethyl acetate (200 ml) was added and the organic phase was washed twice with 5% acetic acid, 5% sodium hydrogencarbonate and water. The organic phase was dried

30

with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography with hexane/ethyl acetate. Yield: 10.2g = 96%.

b) 2-amino-2-methyl-1-(N-benzyloxycarbonyl-L-valyloxy)-propane.

- 5 To a solution of 2-(N-(tert.-butoxycarbonyl)-amino)-2-methyl-1-(N-benzyloxycarbonyl-L-valyloxy)-propane (10g, 23 mmole) in dichloromethane (150 ml) was added trifluoroacetic acid (30 ml) and the mixture was stirred for 1 hour at room temperature. The solution was evaporated under reduced pressure and 10% sodium carbonate solution was added. The product was  
10 extracted four times with dichloromethane, dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography with dichloromethane/methanol. Yield: 3.0g = 40% (crude)

- c) 2-(N-(chloromethoxycarbonyl)-amino)-2-methyl-1-(N-benzyloxycarbonyl-L-valyloxy)-propane.  
15

To a solution of 2-amino-2-methyl-1-(N-benzyloxycarbonyl-L-valyloxy)-propane (2.9g, 9 mmole) and pyridine (2 ml) in dichloromethane (50 ml) was added chloromethyl chloroformate (1.55g, 12 mmole) and the mixture was stirred for 3 hours at room temperature. The mixture was evaporated under  
20 reduced pressure and ethyl acetate was added. The organic phase was washed with water, dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography with hexane/ethyl acetate. Yield: 1.1g = 29%.


- d) 2-(N-(iodomethoxycarbonyl)-amino)-2-methyl-1-(N-benzyloxycarbonyl-L-valyloxy)-propane.  
25

To a solution of 2-(N-(chloromethoxycarbonyl)-amino)-2-methyl-1-(N-benzyloxycarbonyl-L-valyloxy)propane (1.05g, 2.53 mmole) in dry acetone (20 ml) was added sodium iodide (1.8g, 12 mmole) and the mixture was stirred for 36 hours at room temperature. The mixture was evaporated under reduced  
30 pressure and ethyl acetate and water were added. The organic phase was washed with 10% sodium thiosulfate solution and water. The organic phase was dried with sodium sulfate and evaporated under reduced pressure. Yield: 1.04g = 81%.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) 0.92 (m, 6H) 1.35 (s, 6H) 2.10 (m, 1H) 3.88 (m, 1H) 4.35 (m, 2H) 5.11 (s, 2H) 5.32 (d, 1H) 5.82 (s, 1H) 5.91 (s, 2H) 7.35 (m, 5H)

Example A-I-25

- 5 1-(2-N-CBz-L-valyloxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid iodomethyl ester



- a) 6-oxo-1,6-dihydro-pyridine-3-carboxylic acid 4-methoxybenzyl ester.

To a solution of 6-hydroxynicotinic acid (4.87 g, 35 mmol) in DMF (100 mL) at room temperature, was added potassium tert-butoxide (3.93 g, 35 mmol). The reaction mixture was stirred at 60 °C for 1h. 4-Methoxybenzylchloride (8.30 g, 53 mmol) was added and the reaction mixture was stirred at 60 °C for 4h. The DMF was evaporated under vacuum, the residue was dissolved in ether (200 mL) and washed with water (3 x 100 mL). The organic phase was dried with  $\text{Na}_2\text{SO}_4$  and evaporated to give 4.41 g of 6-oxo-1,6-dihydro-pyridine-3-carboxylic acid 4-methoxybenzyl ester.

- b) 1-(2-Hydroxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid 4-methoxybenzyl ester

To a solution of 6-oxo-1,6-dihydro-pyridine-3-carboxylic acid 4-methoxybenzyl ester (4.41 g, 17 mmol) and  $\text{K}_2\text{CO}_3$  (2.58 g, 18.7 mmol) in DMF (100 mL) at room temperature, was added 2-bromoethanol (2.02 g, 16.2 mmol). The reaction mixture was stirred at 80 °C for 30h, whereupon the DMF was evaporated under vacuum. The crude product was column chromatographed (silica gel, 2→5% MeOH in  $\text{CH}_2\text{Cl}_2$ ), to give 3.91 g of 1-(2-hydroxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid 4-methoxybenzyl ester.

- c) 1-(2-N-CBz-L-valyloxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid 4-methoxybenzyl ester

To a mixture of DCC (5.06 g, 24.5 mmol), DMAP (318 mg, 2.6 mmol) and N-CBz-L-valine (6.48 g, 25.8 mmol) in  $\text{CH}_2\text{Cl}_2$  (200 mL) at 0 °C, was added

dropwise a solution of 1-(2-hydroxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid 4-methoxybenzyl ester (6.40 g, 24 mmol) in  $\text{CH}_2\text{Cl}_2$  (200 mL). After 1h at 0 °C, the temperature of the reaction mixture was allowed to assume room temperature and then the mixture was stirred for 5h at room temperature. The mixture was filtered through a glass filter and the solvent was removed under reduced pressure. The crude product was column chromatographed (silica gel, 2→5% MeOH in  $\text{CH}_2\text{Cl}_2$ ), to give 6.81 g 1-(2-N-CBz-L-valyloxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid 4-methoxybenzyl ester.

10 d) 1-(2-N-CBz-L-valyloxyethyl)-2-pyridone-5-carboxylic acid

To a solution of 1-(2-N-CBz-L-valyloxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid 4-methoxybenzyl ester (6.46 g, 12 mmol) in  $\text{CH}_2\text{Cl}_2$  (85 mL) at room temperature, was added trifluoroacetic acid (15 mL). After 1h at room temperature, the reaction mixture was concentrated under reduced pressure.

15 The crude product was column chromatographed (silica gel, 3→6% MeOH in  $\text{CH}_2\text{Cl}_2$ ), to give 4.91 g 1-(2-N-CBz-L-valyloxyethyl)-2-pyridone-5-carboxylic acid.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 12.15 (br s, 1H), 8.29 (d,  $J = 2.2$  Hz, 1H), 7.93 (dd,  $J = 9.5$ , 2.2 Hz, 1H), 7.31 (m, 5H), 6.69 (d,  $J = 9.5$  Hz, 1H), 5.53 (d, 1H), 5.07 (s, 2H),  
20 4.52-4.05 (m, 5H), 2.20-2.00 (m, 1H), 0.90 (d, 3H), 0.81 (d, 3H).

e) 1-(2-N-CBz-L-valyloxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid chloromethyl ester

To a solution of 1-(2-N-CBz-L-valyloxyethyl)-2-pyridone-5-carboxylic acid (4.91 g, 11.8 mmol) in dioxane (200 mL), was added dropwise a 40%

25 aqueous solution of tetrabutylammonium hydroxide (7.65 g). After stirring for 5 min, the solution was evaporated to dryness through co-evaporation with dioxane and toluene. The residue was dissolved in dichloromethane (200 mL) and then chloriodomethane (8.74 mL, 120 mmol) was added and the solution was stirred for 12h at room temperature. The solution was concentrated under  
30 reduced pressure and the residue was shaken with hexane / ethyl acetate (1:1 v/v, 200 mL). The yellow crystalline solid was filtered off and the filtrate was washed with aqueous solution of sodium thiosulfate (0.1 M) and the filtered through anhydrous sodium sulfate and evaporated to dryness. The residue

was column chromatographed (silica gel, 2-4% MeOH in CH<sub>2</sub>Cl<sub>2</sub>), to give 1.80 g of 1-(2-N-CBz-L-valyloxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid chloromethyl ester.


- f) 1-(2-N-CBz-L-valyloxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid iodomethyl ester

To a solution of 1-(2-N-CBz-L-valyloxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid chloromethyl ester (1.80 g, 3.87 mmol) in acetonitrile (30 mL), was added sodium iodide (2.32 g, 15.5 mmol). The solution was stirred for 4 h at 60 °C. The resulting suspension was filtered and the filtrate was evaporated. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and washed with aqueous sodium thiosulfate (0.1 M). The organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure. The crude product was column chromatographed (silica gel, 1% MeOH in CH<sub>2</sub>Cl<sub>2</sub>), to give 2.04 g 1-(2-N-CBz-L-valyloxyethyl)-6-oxo-1,6-dihydro-pyridine-3-carboxylic acid iodomethyl ester.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 8.19 (d, *J* = 2.5 Hz, 1H), 7.79 (dd, *J* = 9.6, 2.5 Hz, 1H), 7.32 (m, 5H), 6.52 (d, *J* = 9.6 Hz, 1H), 6.04 (s, 2H), 5.38 (d, 1H), 5.07 (s, 2H), 4.54-4.06 (m, 5H), 2.20-2.00 (m, 1H), 0.91 (d, 3H), 0.81 (d, 3H).

#### Example A-I-26

##### Iodomethyl 5-[(N-benzyloxycarbonyl-L-valyloxy)methyl]-2-furoate



##### (a) 5-[(N-Benzyloxycarbonyl-L-valyloxy)methyl]-2-furaldehyde

- A solution of 5-(hydroxymethyl)-2-furaldehyde (1.00 g, 7.69 mmol) in 5 mL dry CH<sub>2</sub>Cl<sub>2</sub> was added to a mixture of *N*-benzyloxycarbonyl-L-valine (2.40 g, 9.57 mmol), *N,N'*-dicyclohexylcarbodiimide (2.00 g, 9.69 mmol), and 4-dimethylaminopyridine (117 mg, 0.96 mmol) in 45 mL CH<sub>2</sub>Cl<sub>2</sub>. After stirring overnight, the reaction slurry was filtered, concentrated under vacuum, and subjected to

flash column chromatography (silica, 2/1 petroleum ether - ethyl acetate to give the valine ester (quantitative yield).

(b) 5-[(*N*-Benzyloxycarbonyl-L-valyloxy)methyl]-2-furoic acid

- A solution of NaClO<sub>2</sub> (2.8 mmol) in 3 mL water was added dropwise to a stirred solution of 5-[(*N*-benzyloxycarbonyl-L-valyloxy)methyl]-2-furaldehyde (798 mg, 2.22 mmol) from step (a) in 3 mL MeCN, with cooling in an ice bath. After 2.5 h, the ice bath was removed, 2 mL more MeCN was added, and the two-phase liquid reaction mixture was stirred at room temperature for 25 h. The reaction mixture was diluted with water, made basic with saturated NaHCO<sub>3</sub>, and extracted with ethyl acetate (3 x 50 mL). The separated aqueous solution was acidified to pH 2 with 5% aqueous HCl and extracted with ethyl acetate (3 x 50 mL). This second ethyl acetate solution was washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, and evaporated to dryness under vacuum to give the carboxylic acid (287 mg, 34%) which was used in the next step without further purification.

<sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) 0.84 and 0.93 (2d, 3H each, *J* = 6.8 Hz), 2.15 (m, 1H), 4.35 (dd, 1H, *J* = 9.0, 4.7 Hz), 5.10-5.24 (m, 4H), 5.44 (d, 1H, *J* = 9.0 Hz), 6.54 (d, 1H, *J* = 3.3 Hz), 7.23 (d, 1H, *J* = 3.3 Hz), 7.33 (s, 5H), 11.05 (br s, 1H).

- (c) Chloromethyl 5-[(*N*-benzyloxycarbonyl-L-valyloxy)methyl]-2-furoate Tetrabutylammonium hydroxide (40 wt. % solution in water, 0.55 mL, 0.84 mmol) was added to the carboxylic acid (286 mg, 0.76 mmol) from step (b) in 5 mL dioxane. The yellow solution was concentrated under vacuum, coevaporating several times with dioxane, toluene, and, lastly, CH<sub>2</sub>Cl<sub>2</sub>. The residue was charged with 10 mL dry CH<sub>2</sub>Cl<sub>2</sub> and chloriodomethane (0.55 mL, 7.55 mmol) was added. After stirring for 20.5 h, the reaction mixture was concentrated and subjected to flash column chromatography (silica, 2/1 petroleum ether - ethyl acetate) to give the chloromethyl ester (137 mg, 42%).

- (d) Iodomethyl 5-[(*N*-benzyloxycarbonyl-L-valyloxy)methyl]-2-furoate All of the chloromethyl ester (137 mg, 0.32 mmol) from step c) was refluxed with NaI (195 mg, 1.3 mmol) in 3.2 mL dry MeCN at 70 °C for 4 h. The solvent was removed under vacuum and the residue was subjected to flash column chromatography (silica, 3/1 petroleum ether - ethyl acetate) to give the iodomethyl ester (152 mg, 92%).

<sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) 0.84 and 0.93 (2d, 3H each, *J* = 6.8 Hz), 2.16 (m, 1H), 4.33 (dd, 1H, *J* = 9.1, 4.7 Hz), 5.09-5.21 (m, 4H), 5.36 (d, 1H, *J* = 9.1 Hz), 6.08 (s, 2H), 6.52 (d, 1H, *J* = 3.4 Hz), 7.19 (d, 1H, *J* = 3.5 Hz), 7.33 (s, 5H).

#### Example A-I-27

##### 4-(2-N-benzyloxycarbonyl-L-valyloxyethyl)benzoic acid .

###### a) 4-Methoxybenzyl 4-(2-hydroxyethoxy)benzoate

To a solution of 4-methoxybenzyl 4-hydroxybenzoate (7.0g, 27 mmole) in dry N,N-dimethylformamide (50 ml) was added potassium carbonate (4.15g, 30 mmole) and 2-bromoethanol. The mixture was stirred 48 hours at 80°C, evaporated under reduced pressure and ethyl acetate and water were added. The organic phase was washed five times with water and dried with sodium sulfate. The solution was evaporated under reduced pressure and the product was isolated by silica gel column chromatography with hexane/ethyl acetate. Yield: 6.8g = 83%.

###### b) 4-methoxybenzyl 4-(2-N-benzyloxycarbonyl-L-valyloxyethoxy)benzoate.

To a solution of 4-methoxybenzyl 4-(2-hydroxyethoxy) benzoate (6.6g, 21.8 mmole), 4-dimethylaminopyridine (0.61g, 5 mmole) and N-benzyloxycarbonyl-L-valine (6.3g, 25 mmole) in dichloromethane (80 ml) was added dicyclohexylcarbodiimide (5.2g, 25 mmole) and the mixture was stirred overnight at room temperature. The mixture was cooled and the urethane was filtered. The solution was evaporated and ethyl acetate (200 ml) was added. The organic phase was washed twice with 5% acetic acid, 5% sodium hydrogencarbonate and water. The organic phase was dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography with dichloromethane/methanol. Yield: 10.6g = 90 %.

###### c) 4-(2-N-benzyloxycarbonyl-L-valyloxyethoxy)-benzoic acid.

To a solution of 4-methoxybenzyl 4-(2-N-benzyloxycarbonyl-L-valyloxyethoxy) benzoate (10.2g, 19.04 mmole) in dichloromethane (100 ml) was added trifluoroacetic acid (20 ml) and the mixture was stirred 3 hours at room



temperature. The solution was evaporated under reduced pressure and co-evaporated two times with toluene. The product was isolated by silica gel column chromatography. Yield: 6.9g = 87%. The product may be activated and esterified direct to a drug or converted to iodomethyl 4-(2-N-benzyloxycarbonyl-L-valyloxyethoxy)-benzoic acid as described above, that is by treatment with a base, chloriodomethane, separation and then treatment with NaI.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.94 (m, 6H) 2.18 (m, 1H) 4.22- 4.68 (m, 5H) 5.10 (s, 2H)  
6.94 (d, 2H) 7.35 (m, 5H) 8.05 (d, 2H)

#### Preparation of compounds of the invention

##### Example 1

(1S, 2S)-N-{*cis*-2-[6-fluoro-2-(L-isoleucyloxymethyloxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea

a) (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(N-BOC-L-isoleucyloxymethyloxy)-3-propionylphenyl] cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea.

To a solution of (1S, 2S)-N-{*cis*-2-[6-fluoro-2-hydroxy-3-propionylphenyl] cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea (2.03 g, 5.5 mmol) in THF (50 mL) at 20 °C, was added NaH (60%, 220 mg, 5.5 mmol). After the mixture was stirred 1.5h at 20 °C, N-BOC-L-isoleucine iodomethyl ester (16.5 g, 16.5 mmol) was added. The solution was stirred for 6h at room temperature and then concentrated under reduced pressure. The crude product was column chromatographed (aluminium oxide 90, 1% MeOH in CH<sub>2</sub>Cl<sub>2</sub>), to give 1.76 g of the title product.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 9.75 (br s, 1H), 9.15 (br s, 1H), 8.16 (s, 1H), 7.71 (dd, 1H), 7.52 (dd, 1H), 7.00-6.87 (m, 2H), 5.81 (d, 1H), 5.68 (d, 1H), 5.00 (d, 1H), 4.21 (dd, 1H), 3.40-3.25 (m, 1H), 2.99-2.72 (m, 2H), 2.10 (dd, 1H), 1.85-1.68 (m, 1H), 1.60-1.47 (m, 1H), 1.41 (s, 9H), 1.32-1.05 (m, 3H), 1.13 (t, 3H), 0.88-0.78 (m, 6H).

b) (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(L-isoleucyloxymethoxy)-3-propionylphenyl] cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea

To TFA (30 mL) at 0 °C, was added (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(N-BOC-L-isoleucyloxymethoxy)-3-propionylphenyl] cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea (1.81 g, 2.96 mmol). The reaction mixture was stirred at 0 °C for 30 min and then concentrated under reduced pressure at 0 °C. The crude product was column chromatographed (silica gel, 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub>), to give 1.48 g of the title compound as the TFA-salt.  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 9.50 (br s, 1H), 9.42 (br s, 1H), 8.34 (s, 1H), 7.73 (dd, 1H), 7.27 (m, 1H), 7.10 (d, 1H), 6.81 (dd, 1H), 6.16 (d, 1H), 5.73 (d, 1H), 3.87 (d, 1H), 3.39 (m, 1H), 3.05-2.68 (m, 2H), 2.29 (dd, 1H), 2.10-1.88 (m, 2H), 1.57-1.21 (m, 3H), 1.09 (t, 3H), 1.02 (d, 3H), 0.91 (t, 3H).

#### Example 2

15 (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(L-valyloxymethoxy)-3-propionylphenyl]} cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea

a) (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(N-CBz-L-valyloxymethoxy)-3-propionylphenyl]}cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea .

20 To a solution of (1S, 2S)-N-{*cis*-2-[6-fluoro-2-hydroxy-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea (368 mg, 1 mmole ) in THF (5 ml) was added sodium hydride in paraffin (60 %, 38 mg, 0.95 mmole). After 1.5 hour, N-CBz-L-valyloxymethyl iodide (1.09g, 2.8 mmole) prepared analogously to the N-BOC-L-isoleucyloxymethyl iodide described  
25 above was added to the solution and reaction was kept 18 hours. The mixture was filtered through Celite and poured into sodium hydrogen carbonate aqueous solution, and extracted with methylene chloride. The organic phase was dried and the product was isolated with silica gel column chromatography to yield 210 mg.  
30 <sup>1</sup>H-NMR (CDCl<sub>3</sub>): 8.16 (s, 1H), 7.70 (dd, 1H), 7.49 (t, 1H), 7.35 (m, 5H), 6.93 (m, 2H), 5.78 (dd, 2H), 5.27 (d, 1H), 5.11 (s, 2H), 4.28 (m, 1H), 3.34 (m, 1H), 2.84 (m, 2H), 2.09 (m, 2H), 1.54 (m, 1H), 1.34 (m, 1H), 1.10 (t, 3H), 0.87 (dd, 6H).

b) (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(L-valyloxymethoxy)-3-propionylphenyl]]cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea

(1S, 2S)-N-{*cis*-2-[6-fluoro-2-(N-CBz-L-valyloxymethoxy)-3-propionylphenyl]]cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea (200 mg, 0.32

5 mmole) was dissolved in a mixed solvent of methanol (5 ml), ethylacetate (2 ml) and acetic acid (1 ml). To the solution was added palladium black (35 mg). It was kept under hydrogen at atmospheric pressure for two hours. After filtration, the solution was evaporated and the product was purified by silica gel column chromatography yielding 66 mg.

10

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 8.20 (d, 1H), 7.73 (dd, 1H), 7.44 (dd, 1H), 6.94 (m, 2H), 5.80 (dd, 2H), 3.37 (1H), 2.88 (m, 2H), 2.10 (m, 2H), 1.60 (m, 1H), 1.46 (m, 1H), 1.08 (t, 3H), 0.94 (m, 6H).

15 Example 3

(1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2,2-dimethyl-3-(L-valyloxy)-propionyloxy-methoxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea

a) (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2,2-dimethyl-3-(N-Boc-L-

20 valyloxy)propionyloxymethoxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea.

To a solution of (1S, 2S)-N-[*cis*-2-(6-fluoro-2-hydroxy-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)] urea (368 mg, 1 mmole) in THF (5 ml)

was added sodium hydride in paraffin (60 %, 38 mg, 0.95 mmole). After one  
25 hour, 2,2-dimethyl-3-(N-Boc-L-valyloxy)propionic acid iodomethyl ester (1.35g, 3 mmole) was added to the solution. After 5 hr at room temperature, it was then raised to 50 °C and reaction was kept 18 hours. The reaction mixture was poured into sodium hydrogen carbonate aqueous solution and extracted with methylene chloride. The organic phase was dried and the  
30 product was isolated with alumina column chromatography. 140 mg.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 8.39-6.70 (m, 5H) 5.77 (m, 2H) 5.15 (d, 1H) 4.00 (m, 3H) 3.40 (m, 1H) 2.90 (m, 2H) 2.30 (m, 1H) 2.20 (m, 1H) 1.70 (m, 1H) 1.42 (s, 9H) 1.16 (d, 6H) 0.92 (m, 9H)

- b) (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2,2-dimethyl-3-(L-valyloxy)propionyl-oxymethoxy)-3-propionylphenyl]-cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2,2-dimethyl-3-(N-Boc-L-valyloxy)-propionyloxymethoxy)-3-propionylphenyl]} cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea (120 mg) was treated with trifluoroacetic acid at 0° C for 20 min. The solution was evaporated and coevaporated with toluene and methanol succesively, giving the titled product in quantitative yield.
- <sup>1</sup>H-NMR (CDCl<sub>3</sub>): 8.33 (d, 1H) 7.89 (d, 1H) 7.48 (t, 1H) 7.16 (m, 1H) 6.96 (t, 1H) 5.70 (dd, 2H) 4.18 (dd, 2H) 4.01 (m, 1H) 3.38 (m, 1H) 2.88 (m, 2H) 2.16 (m, 1H) 1.58 (m, 2H) 1.25 (d, 6H) 1.04 (m, 9H).

#### Example 4

(1S, 2S)-N-{*cis*-2-[6-fluoro-2-(3,3-bis-(L-valyloxymethyl)propionyloxy-methoxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea

15

- a) (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(3,3-bis ( N-CBz-L-valyloxymethyl) propionyloxymethoxy)-3-propionylphenyl]}cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea .

To a solution of (1S, 2S)-N-{*cis*-2-[6-fluoro-2-hydroxy-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea (331 mg, 1 mmole) in THF (5 ml) was added sodium hydride in paraffin (60 %, 32 mg, 0.81 mmole). After one hour, 3,3-bis-(N-CBz-L-valyloxymethyl) propionic acid iodomethyl ester (1.3g, 1.8 mmole) was added to the solution. After 5 hr at room temperature, it was then raised to 50 °C and reaction was kept 18 hours. The mixture was poured into sodium hydrogen carbonate aqueous solution, and extracted with methylene chloride. The organic phase was dried and the product was isolated with alumina column chromatography. 185 mg.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 8.19 (s, 1H) 7.89 (dd, 1H) 7.32 (m, 11H) 7.10 (m, 1H) 6.90 (t, 1H) 5.79 (dd, 2H) 5.09 (s, 2H) 4.31 (m, 2H) 4.08 (m, 4H) 2.95 (m, 2H) 2.50 (m, 3H) 2.17 (m, 3H) 1.55 (m, 1H) 1.07 (t, 3H) 0.88 (dd, 12 H).

- b) (1S, 2S)-N-{*cis*-2-[6-fluoro-2-( 3,3-bis (L-valyloxymethyl) propionyloxy-methoxy)-3-propionylphenyl]}cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea.

(1S, 2S)-N-{*cis*-2-[6-fluoro-2-(3,3-bis ( N-CBz-L-valyloxymethyl) propionyloxymethoxy )-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea (170 mg, 0.17 mmole) was dissolved in a mixed solvent of methanol (5 ml), ethyl acetate (2 ml) and acetic acid (1 ml). To the solution  
5 was added palladium black (30 mg). It was kept under hydrogen at atmospheric pressure for four hours. After filtration, the solution was evaporated and the product was purified by silica gel column chromatography. 80 mg.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 8.38 (d, 1H) 8.02 (d, 1H) 7.42 (m, 2H) 7.12 (t, 1H) 5.70  
10 (dd, 2H) 4.00 (s, 4H) 3.16 (m, 1H) 3.08 (d, 2H) 2.80 (m, 1H) 2.40 (m, 2H), 2.11 (m, 1H) 1.52 (m, 1H) 0.95 (t, 3H) 0.98 (dd, 12 H).

#### Example 5

(1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2-(L-valyloxy)-ethoxycarbonyloxymethoxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea  
15

a) (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2-(N-CBz-L-valyloxy)-ethoxycarbonyloxymethoxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea.

20 To a solution of (1S, 2S)-N-{*cis*-2 [6-fluoro-2-hydroxy-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea (368 mg, 1 mmole) in THF (5 ml) was added sodium hydride in paraffin (60 %, 38 mg, 0.95 mmole). After 1.5 hr, 2- (N-CBz-L-valyloxy)ethoxycarbonyloxymethyl iodide (864 mg, 1.7 mmole) was added to the solution. The reaction was kept for 48 hours. The  
25 mixture was poured into sodium hydrogen carbonate aqueous solution, and extracted with methylene chloride. The organic phase was dried and the product was isolated with silica gel column chromatography. 210 mg.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 8.21 (d, 1H) 7.72 (d, 1H) 7.28 (m, 6H) 6.90 (m, 2H) 5.75 (dd, 2H) 5.09 (s, 2H) 4.35 (m, 4H) 2.85 (m, 2H) 2.50 (m, 2H) 2.16 (m, 1H),  
30 1.65 (m, 1H) 1.11 (t, 3H) 0.93 (dd, 6 H).

b) 1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2-(L-valyloxy)-ethoxycarbonyloxymethoxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea.

(1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2-(N-CBz-L-valyloxy)-ethoxycarbonyloxymethyloxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea is deprotected by conventional techniques such as palladium black in a mixed solvent of methanol, ethyl acetate and acetic acid under hydrogen at atmospheric pressure followed by conventional work up such as filtration, evaporation and silica gel column chromatography.

#### Example 6

(1S,2S)-N-[*cis*-2-(6-fluoro-2-(1,3-bis-L-valyloxy-2-(propoxycarbonyloxymethyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea

a) (1S,2S)-N-[*cis*-2-(6-fluoro-2-(1,3-bis-(N-BOC-L-valyloxy-2-(propoxycarbonyloxymethyloxy)-3-propionylphenyl)cyclopropyl)-N'-[2-(5-cyanopyridyl)]urea.

NaH (121 mg, 60% w/w in mineral oil, 3.0 mmol) was added to a mixture of (1S,2S)-N-[*cis*-2-(6-fluoro-2-hydroxy-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea (1.05 g, 2.85 mmol) in 15 mL dry THF under N<sub>2</sub>. After 1 h, the solution was concentrated to dryness and redissolved in 10 mL DMF. 2-O-iodomethoxycarbonyl-1,3-di-O-(*N*-tert-butoxycarbonyl-L-valyl)glycerol (2.96 g, 4.39 mmol) in 15 mL DMF was added and the reaction mixture was stirred for 20 h. Removal of solvent under vacuum followed by flash column chromatography (silica gel, 2/1 ethyl acetate - petroleum ether) gave 1.46 g (56%) of the title product as a white solid.

<sup>1</sup>H NMR (250 MHz, CD<sub>3</sub>OD) 0.94 and 0.97 (2d overlap, 12H), 1.11 (t, 3H), 1.23 (m, 1H), 1.46 (s, 18H), 1.64 (m, 1H), 2.07-2.24 (m, 3H), 2.90 (m, 2H), 3.32 (m, 1H), 4.06 (d, 2H), 4.28-4.52 (m, 4H), 5.13 (m, 1H), 5.78 and 5.88 (AB q, 2H), 7.07-7.19 (m, 2H), 7.62 (dd, 1H), 7.92 (dd, 1H), 8.31 (d, 1H).

b) (1S,2S)-N-[*cis*-2-(6-fluoro-2-(1,3-bis-L-valyloxy-2-(propoxycarbonyloxymethyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea.

Ice-cold trifluoroacetic acid (30 mL) was added to the intermediate of step a (1.69 g, 1.85 mmol) in an ice bath, under N<sub>2</sub>. After 7 min, the reaction mixture was concentrated under vacuum, coevaporating several times with, initially,

toluene and, finally, CH<sub>2</sub>Cl<sub>2</sub>. The oily residue was chromatographed immediately on a silica gel column with 10-20 % methanol in CH<sub>2</sub>Cl<sub>2</sub> to give 1.37 g of the product as a trifluoroacetate salt.

5 <sup>1</sup>H NMR (250 MHz, CD<sub>3</sub>OD) 1.07-1.12 (m, 15H), 1.26 (m, 1H), 1.63 (m, 1H), 2.19 (m, 1H), 2.35 (m, 2H), 2.89 (m, 2H), 4.08 (m, 2H), 4.44-4.71 (m, 4H), 5.26 (m, 1H), 5.79 and 5.91 (AB q, 2H), 7.10-7.18 (m, 2H), 7.59 (dd, 1H), 7.93 (dd, 1H), 8.30 (d, 1H).

<sup>19</sup>F NMR (235 MHz, CD<sub>3</sub>OD) -103.5, -73.5.

10

#### Example 7

(1S,2S)-N-[cis-2-(6-fluoro-2-(L-valyloxy)methoxycarbonyloxy-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea

15 a) (1S,2S)-N-[cis-2-(6-fluoro-2-chloromethoxycarbonyloxy-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea.

Chloromethyl chloroformate (2.3 mL, 25 mmol) was added by syringe to a mixture of (1S,2S)-N-[cis-2-(6-fluoro-2-hydroxy-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea (4.695 g, 12.7 mmol) and pyridine  
20 (6.1 mL, 76 mmol) in 65 mL dry CH<sub>2</sub>Cl<sub>2</sub> with cooling in an ice bath, under N<sub>2</sub>. After 10 min, the ice bath was removed and the mixture was stirred at room temperature for 1h 40 min. The mixture was diluted with 100 mL CH<sub>2</sub>Cl<sub>2</sub> and washed with 50 mL H<sub>2</sub>O. The aqueous phase was reextracted with 25 mL CH<sub>2</sub>Cl<sub>2</sub>. The combined organic phases were washed with 50 mL saturated  
25 NaHCO<sub>3</sub>, followed by 2 x 50 mL brine. Drying over Na<sub>2</sub>SO<sub>4</sub> and concentration under vacuum gave a crude material that was subjected to flash column chromatography (silica gel, 1/1 ethyl acetate - petroleum ether) to give 4.05 g (69%) title product.

<sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) 1.15 (t, 3H), 1.30 (m, 1H), 1.59 (m, 1H), 2.02  
30 (m, 1H), 2.87 (q, 2H), 3.29 (m, 1H), 5.87 (s, 2H), 6.97 (d, 1H), 7.09 (m, 1H), 7.72 (dd, 1H), 7.76 (dd, 1H), 8.10 (dd, 1H), 9.26 (br s, 1H), 10.09 (brs, 1H).

b) (1S,2S)-N-[*cis*-2-(6-fluoro-2-iodomethoxycarbonyloxy-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea

(1S,2S)-N-[*cis*-2-(6-fluoro-2-chloromethoxycarbonyloxy-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea (3.97 g, 8.6 mmol) and NaI (5.17 g, 34.5 mmol) in 85 mL dry acetonitrile were refluxed at 70 °C for 4 h under N<sub>2</sub>. The solvent was removed *in vacuo*, the residue was partitioned between 100 mL CH<sub>2</sub>Cl<sub>2</sub> and 25 mL H<sub>2</sub>O, the aqueous phase was reextracted with 10 mL CH<sub>2</sub>Cl<sub>2</sub>, and the organic phases were combined, washed successively with 2 x 25 mL 5% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> and 2 x 25 mL brine, and dried over Na<sub>2</sub>SO<sub>4</sub>. Flash column chromatography (silica gel, 2/1 ethyl acetate - petroleum ether) of the crude product obtained after concentration *in vacuo* gave 4.15 g material containing 92% of the title compound and traces of the starting material. <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) 1.18 (t, 3H), 1.34 (m, 1H), 1.62 (m, 1H), 2.03 (m, 1H), 2.86 (q, 2H), 3.32 (m, 1H), 6.08 (s, 2H), 6.97 (d, 1H), 7.08 (m, 1H), 7.70-7.76 (m, 2H), 8.13 (d, 1H), 8.90 (br s, 1H), 9.30 (br s, 1H).

c) (1S,2S)-N-[*cis*-2-(6-fluoro-2-(N-BOC-L-valyloxy)methoxycarbonyloxy-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea

Tetrabutylammonium hydroxide (40 wt % solution in water, 6.4 mL, 9.8 mmol) was added to Boc-L-valine (2.54 g, 11.7 mmol) in 30 mL dioxane. The solution was concentrated *in vacuo*, coevaporating several times with dioxane, toluene, and CH<sub>2</sub>Cl<sub>2</sub>, and dried under vacuum overnight. The resulting Q salt was dissolved in 30 mL dry CH<sub>2</sub>Cl<sub>2</sub> and (1S,2S)-N-[*cis*-2-(6-fluoro-2-(iodomethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea (7.1 mmol) in 65 mL dry CH<sub>2</sub>Cl<sub>2</sub> was added. After stirring under N<sub>2</sub> for 18 h, the reaction mixture was washed with 3 x 50 mL H<sub>2</sub>O, 1 x 50 mL 5% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, and 2 x 50 mL H<sub>2</sub>O. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub>, concentrated, and submitted to flash column chromatography (silica gel, 3/1 ethyl acetate - petroleum ether) to give 2.21 g (49%) product. <sup>1</sup>H NMR (250 MHz, CD<sub>3</sub>OD) 0.98 (d, 3H), 1.02 (d, 3H), 1.17 (t, 3H), 1.24 (m, 1H), 1.47 (s, 9H), 1.59 (m, 1H), 2.06 (m, 1H), 2.24 (m, 1H), 2.96 (q, 2H), 3.24 (m, 1H), 4.15 (d, 1H), 5.94 and 6.02 (AB q, 2H), 7.12 (d, 1H), 7.26 (m, 1H), 7.91 (dd, 1H), 7.94 (dd, 1H), 8.23 (dd, 1H).



d) (1S,2S)-N-[*cis*-2-(6-fluoro-2-(L-valyloxy)methoxycarbonyloxy-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea

Cold trifluoroacetic acid (40 mL) was added to (1S,2S)-N-[*cis*-2-(6-fluoro-2-(N-BOC-L-valyloxymethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea (1.94 g, 3.02 mmol) with cooling in an ice bath, under N<sub>2</sub>. After 5 min, the solution was concentrated *in vacuo*, coevaporating several times with toluene, and then CH<sub>2</sub>Cl<sub>2</sub>, and dried under vacuum for several hours to give the compound as a trifluoroacetate salt in quantitative yield.

<sup>1</sup>H NMR (250 MHz, CD<sub>3</sub>OD) δ 1.12-1.18 (m, 9H), 1.25 (m, 1H), 1.59 (m, 1H), 2.07 (m, 1H), 2.47 (m, 1H), 2.97 (q, 2H), 3.26 (m, 1H), 4.16 (d, 1H), 6.01 and 6.37 (AB q, 2H), 7.11 (d, 1H), 7.29 (m, 1H), 7.92 (dd, 1H), 7.99 (dd, 1H), 8.22 (d, 1H).

<sup>19</sup>F NMR (235 MHz, CD<sub>3</sub>OD) δ -102.7, -74.0.

15

#### Example 8

(1S, 2S)-N-{*cis*-2-[6-fluoro-2-(3-carboxypropionyloxymethyloxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea

a) (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(3-benzyloxycarbonylpropionyl-oxymethyloxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)]urea. 3-Benzyloxycarbonylpropionic acid iodomethyl ester (522 mg, 1.5 mmole) was added to a solution of (1S, 2S)-N-{*cis*-2-[6-fluoro-2-hydroxy-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea (185 mg, 0.5 mmole) in THF (5 ml) which had been treated with sodium hydride in paraffin (60 %, 20 mg, 0.5 mmole) for 30 min. After 18 hr at room temperature, the reaction mixture was poured into sodium hydrogen carbonate aqueous solution, and extracted with methylene chloride. The organic phase was dried and the product was isolated with alumina column chromatography. 115 mg.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 8.20 (d, 1H) 7.72 (dd, 1H) 7.49 (dd, 1H) 7.35 (m, 5H) 6.97 (m, 2H) 5.73 (dd, 2H) 5.17 (s, 2H) 3.35 (m, 1H) 2.88 (m, 2H) 2.60 (m, 4H) 2.12 (m, 1H) 1.58 (m, 1H) 1.11 (t, 3H).

b) (1S, 2S)-N-{*cis*-2-[6-fluoro-2-(3-carboxylpropionyloxymethoxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea.

(1S, 2S)-N-{*cis*-2-[6-fluoro-2-(3-carboxylpropionyloxymethoxy)-3-propionylphenyl]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea (100 mg, 0.17 mmole) was dissolved in a mixed solvent of ethylacetate (3 ml) and acetic acid (1 ml). To the solution was added palladium black (30 mg). It was kept under hydrogen at atmospheric pressure for three hours. After filtration, the solution was evaporated and the product was purified by silica gel column chromatography. 81 mg. An Rx bearing intermediate linking group such as 1,3-bis-N-Boc-valyloxyglycerol (preparation described in WO9909031) can be esterified to the free carboxyl group using conventional esterification conditions as described in WO9909031.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 8.21 (s, 1H) 7.75 (d, 1H) 7.49 (dd, 1H) 7.08 (d, 5H) 6.97 (t, 1H) 5.73 (dd, 2H) 5.17 (s, 2H) 3.26 (m, 1H) 2.87 (m, 2H) 2.60 (m, 4H) 2.09 (m, 1H) 1.58 (m, 1H) 1.11 (t, 3H)

#### EXAMPLE 9

(1S, 2S)-N-[*cis*-2-(6-fluoro-2-O-(4-L-valyloxybenzoyl)-3propionylphenyl)-cyclopropyl]-N'-(5-cyanopyrid-2-yl) urea

20

a) 4-benzyloxybenzoic acid.

To a solution of 4-hydroxybenzoic acid (6.9g, 50 mmole) in 150 ml DMF was added potassium tert.-butoxide (12.34g, 110 mmole) and the mixture was stirred at room temperature for one hour. Benzyl bromide (20.5g, 120 mmole) was added and the mixture was stirred for two days at room temperature. The mixture was evaporated under reduced pressure and 100ml 1,4-dioxane and a solution of sodium hydroxide (6.0g, 150 mmole) in 50 ml water was added. The mixture was refluxed for two hours, cooled and evaporated under reduced pressure. Water was added and the mixture was acidified with acetic acid. The product was filtered, washed with cold water and dried. Yield: 10.2g = 89%.

30

b) 4-benzyloxybenzoyl chloride.

To a mixture of 4-benzyloxybenzoic acid (2.28g, 10 mmole) in 20 ml dried dichloromethane were added five drops of DMF and 2.5 ml thionyl chloride.

The mixture was refluxed for three hours and evaporated under reduced

5 pressure. Yield: 2.45g = 100%

c) (1S, 2H)-N-[*cis*-2-(6-fluoro-2-O-(4-benzyloxybenzoyl)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyrid-2-yl) urea .

To a solution of (1S, 2S)-N-[*cis*-2-(6-fluoro-2-hydroxy-3-propionylphenyl)cyclopropyl]-N'-(5-cyanopyrid-2-yl) urea (184mg, 0.5 mmole) in 3 ml DMF was

10 added potassium tert. butoxide (78.5mg, 0.7 mmole) and the mixture was stirred for one hour at room temperature. A solution of 4-

benzyloxybenzoylchloride (185mg, 0.75 mmole) in 1ml DMF was added and the mixture was stirred overnight at room temperature. 40 ml ethyl acetate were added and the organic phase was washed four times with water. The

15 solution was dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography. Yield: 180mg = 62%.

<sup>1</sup>H-NMR (DMSO  $\delta$ -6) 0.92 (m, 4H) 1.31(m, 1H) 1.85 (m, 1H) 2.82 (m, 2H) 3.06 (m, 1H) 5.26 (s, 2H) 7.20 (m 2H) 7.38-8.12 (m, 11H) 8.38 (m, 1H)

20 d) (1S, 2S)-N-[*cis*-2-(6-fluoro-2-O (4-hydroxybenzoyl)-3-propionylphenyl)cyclopropyl]-N'-(5-cyanopyrid-2-yl)] urea-O-4-hydroxybenzoate

A solution of the intermediate of step c) (170 mg, 0.29 mmole) in 15 ml ethyl acetate and 15 ml methanol was hydrogenated with 10% palladium on

charcoal (30mg) three times at room temperature and normal pressure. The

25 catalyst was filtered and washed with ethyl acetate and methanol and the solution was evaporated under reduced pressure. The product was isolated by silica gel column chromatography. Yield: 100 mg = 70%.

<sup>1</sup>H-NMR (DMSO  $\delta$ -6) 0.93 (m, 4H) 1.32 (m, 1H) 1.88 (m, 1H) 2.85 (m, 2H) 3.05 (m, 1H) 6.92 (m, 2H) 7.38 (m, 2H) 8.00 (m, 4H) 8.38 (m, 1H)

30 e) (1S, 2S)-N-[*cis*-2-(6-fluoro-2-O (4-L-valyloxybenzoyl)-3-propionylphenyl)-cyclopropyl]-N'-(5-cyanopyrid-2-yl) urea

An R<sub>2</sub> group, such as N-protected L-valyl is acylated to the exposed ring hydroxy group using conventional acylation conditions as described herein and deprotected to yield a compound of the invention.

5 Example 10

(1S, 2S)-N-[cis-2-(6-fluoro-2-O ((4-iso-leucyloxybenzoyloxymethyl)-3-propionylphenyl)-cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea-O-methylene-4-hydroxybenzoate-O-L-iso-leucyl ester

10 a) Methyl-4-(4-methoxybenzyloxy) benzoate.

To a solution of methyl 4-hydroxybenzoate (6.85g, 45 mmole) in 80 ml DMF was added potassium tert. butoxide (5.6 g, 51 mmole) and the mixture was stirred at room temperature for one hour. 4-Methoxybenzyl chloride (8.3 g, 52 mmole) was added and the mixture was stirred overnight at room

15 temperature. The mixture was evaporated under reduced pressure and 200 ml ethyl acetate was added. The organic phase was washed four times with water, dried with sodium sulfate and evaporated under reduced pressure.

Yield: 12.3g = 100%

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 3.82 (s, 3H) 3.88 (s, 3H) 5.03 (s, 2H) 6.96 (m, 4H) 7.36 (d, 20 2H) 7.98 (d, 2H)

b) 4-(4-methoxybenzyloxy) benzoic acid

To a solution of methyl 4-(4-methoxybenzyloxy) benzoate (12.2 g, 44.8 mmole) in 50 ml 1,4-dioxane was added a solution of lithium hydroxide ( 2.15 g, 89,6 mmole) and the mixture was stirred overnight at 60°C. The mixture

25 was evaporated under reduced pressure and 5% acetic acid was added. The product was filtered, washed with water and dried. Yield: 10.1g = 87%

<sup>1</sup>H-NMR (DMSO δ-6) 3.74 (s, 3H) 5.08 (s, 2H) 6.92 (d, 2H) 7.06 (d, 2H) 7.36 (d, 2H) 7.90 (d, 2H)

c) Chloromethyl 4-(4-methoxybenzyloxy)benzoate

30 To a solution of 4-(4-methoxybenzyloxy) benzoic acid (5.16 g, 20 mmole) in 100 ml 1,4-dioxane was added a 40% solution of tetrabutylammonium hydroxide (14.27 g, 22 mmole) and the mixture was stirred 2 hours at room temperature. The mixture was evaporated under reduced pressure and co-

evaporated two times with 1,4-dioxane and two times with toluene. The dried product was dissolved in 60 ml dichloromethane and iodochloromethane (35.3 g 200 mmole) was added. The solution was stirred for two days at room temperature and evaporated under reduced pressure. About 100 ml ethyl  
5 acetate was added and the organic phase washed twice with water, dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography. Yield: 4.48 g = 73%

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 3.83 (s, 3H) 5.06 (s, 2H) 5.94 (s, 2H) 7.00 (m, 4H) 7.36 (d, 2H) 8.05 (d, 2H)

10 d) Iodomethyl 4-(4-methoxybenzyloxy) benzoate

To a solution of chloromethyl 4-(4-methoxybenzyloxy) benzoate (0.77g, 2.5 mmole) in 15 ml dry acetone was added sodium iodide (1.87g, 12.5 mmole) and the mixture was stirred overnight at room temperature. The mixture was evaporated under reduced pressure and extracted with ethyl acetate/water.

15 The organic phase was washed with a 5% sodium thiosulfate solution, dried with sodium sulfate and evaporated under reduced pressure. Yield 0.86g = 86%

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 3.84 (s, 3H) 5.05 (s, 2H) 6.14 (s, 2H) 6.98 (m, 4H) 7.36 (d, 2H) 8.00 (d, 2H)

20 e) (1S, 2S)-N-[*cis*-2-(6-fluoro-2-O-(4(4-methoxybenzyloxy)-benzoyloxymethyl)-3-propionylphenyl (cyclopropyl)-N'-[2-(5-cyanopyridyl)]urea.

To a solution of (1S, 2S)-N-[*cis*-2-(6-fluoro-2-hydroxy-3-propionylphenyl) cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea (368mg, 1 mmole) in 5 ml DMF was  
25 added a suspension of 60% sodium hydride in mineral oil (44mg, 1.1 mmole) and the mixture was stirred for one hour at room temperature. A solution of iodomethyl-4-(4-methoxybenzyloxy) benzoate (0.84 g, 2.1 mmole) in 2 ml THF was added and the mixture was stirred overnight at room temperature. 50 ml ethyl acetate were added and the organic phase was washed four times with  
30 water, dried with sodium sulfate and evaporated under reduced pressure. The product was isolated by silica gel column chromatography. Yield: 525 mg = 82%

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) 0.91 (m, 3H) 1.32 (m, 1H) 1.60 (m, 1H) 2.04 (m, 1H) 2.90 (m, 2H) 3.20 (m, 1H) 3.82 (s, 3H) 5.04 (s, 2H) 5.84-6.06 (m, 2H) 6.91-8.18 (m, 13H)

f) (1S, 2S)-N-[*cis*-2-(6-fluoro-2-O (4-hydroxybenzoyloxymethyl)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea-O-methylene-4-hydroxybenzoate

To a solution of the intermediate of step e) (100 mg, 0.156 mmole) in 4 ml dichloromethane was added TFA (0.5 ml) and the solution was stirred for one hour at room temperature. The solution was evaporated under reduced pressure and the product was isolated by silica gel column chromatography. Yield: 45mg = 55%.

<sup>1</sup>H-NMR (DMSO  $\delta$ -6) 0.84 (m, 3H) 1.10 (m, 1H) 1.48 (m, 1H) 2.12 (m, 1H) 2.80 (m, 2H) 3.19 (m, 1H) 5.85-6.02 (m, 2H) 6.84 (m, 2H) 7.18 (m, 1H) 7.46 (m, 2H) 7.74 (m, 2H) 8.04 (m, 2H) 8.38 (m, 1H)

g) (1S, 2S)-N-[*cis*-2-(6-fluoro-2-O (4-*isoleucyloxy*benzoyloxymethyl)-3-propionylphenyl)-cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea-O-methylene-4-hydroxybenzoate-O-L-*isoleucyl* ester

An R<sub>2</sub> group, such as N-protected L-*isoleucine* is acylated to the exposed hydroxy group using conventional acylation conditions as described herein and deprotected to yield a compound of the invention.

#### Biological Example 1

##### Bioavailability

The release of a compound of Formula P-2 from orally administered prodrugs of Formulae P3 to P8 were monitored in rats. The compounds of Examples 1 to 6 were made up in a propylene glycol vehicle and orally administered to paired fasted male Sprague Dawley rats at a dose corresponding to 0.027 mmol/kg. At 30, 60, 120, 240 & 360 minutes, 0.2 ml blood were collected, centrifuged and frozen for later analysis. The released drug of Formula P-2, (1S, 2S)-N-[*cis*-2-(6-fluoro-2-hydroxy-3-propionylphenyl) cyclopropyl]-N'-[2-(5-cyanopyridyl)] urea was assayed by HPLC. Aliquots comprising 40-100  $\mu$ l of each plasma sample are mixed with an equal volume of acetonitrile (10

seconds, Vibrofex). The sample is centrifuged (2 min, 14000 RPM) and 30  $\mu$ l of the supernatant is injected into an HPLC system, as follows.

- Pre column: RP-18, 7  $\mu$ m, 15 x 3.2 mm  
Column: YMC basic, 3 $\mu$ m, 150 x 3 mm  
5 Mobile phase: 60 % acetonitrile in 3 mM ammonium acetate, pH 6.4  
Flow rate: 0.4 ml/min  
Detection: UV, 250 nm

TABLE P-1

Example	Bioavailability <sub>0-6 hours</sub>	C <sub>max</sub> $\mu$ M
1	34 %	0.78
2	18 %	0.51
3	27 %	0.64
4	18 %	0.43
6	50 %	1.06
7	70 %	1.5

10


The above bioavailabilities correspond to sustained plasma levels of the active metabolite well above the ED<sub>50</sub> for HIV-1. The HIV-1 activity of the mother compound of formula P2 is described and quantitated in PCT/SE99/00194.

15

Although the invention has been illustrated by reference to examples employing particular mother NNRTIs, particular amino acids and particular linking groups it will be appreciated that the invention is not limited to these values but extends throughout the spirit and scope of the following claims.

Claims

1 A compound of the formula P1:



P-1

wherein:

Ar1 is an unsaturated, optionally substituted, mono or bicyclic ring structure comprising 0 to 3 hetero atoms selected from S, O and N;

10 Ar2 is an aromatic, optionally substituted, monocyclic ring structure comprising at least one nitrogen hetero atom and zero to two further hetero atoms selected from S, O and N;

R4 and R5 are independently H or C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl, C<sub>1</sub>-C<sub>5</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> alkanoyloxy, C<sub>1</sub>-C<sub>4</sub> alkylthio, amino, carboxy, carbamoyl, cyano, halo, hydroxy, aminomethyl, hydroxymethyl, carboxymethyl, or halo substituted C<sub>1</sub>-C<sub>6</sub> alkyl mercapto, nitro; or R4 and R5 join to form a 3 – 6 membered, optionally substituted ring structure;

R6 is O or S;

20 L\* is a linker moiety which is ether-, carbonate- or ester-bound to the adjacent oxygen and ester linked to Rx;


Rx is the residue of a natural or unnatural amino acid;

and pharmaceutically acceptable salts thereof.

25 2. A compound according to claim 1, wherein L\* comprises a C<sub>1</sub>-C<sub>8</sub> alkyloxy group with an ether linkage to said hydroxy residue.



3. A compound according to claim 2, wherein the alkyloxy group has the formula:



5 wherein the arrowed oxygen together with PETT is the hydroxy residue of formula I, Ry and Ry' are independently H or C<sub>1</sub>-C<sub>3</sub> alkyl and A is the Rx residue or an ester or carbonate bonded intermediate linking moiety to which Rx is esterified.

10 4 A compound according to claim 3, wherein Ry and Ry' are both hydrogen.

5 A compound according to claim 2 wherein Rx is esterified to the alkyloxy group.

15

6 A compound according to claim 2 wherein L\* further comprises an intermediate linking moiety esterified or carbonate bonded to the alkyloxy group, wherein the intermediate linking moiety possesses an hydroxy group to which Rx is esterified.


20

7 A compound according to claim 1, wherein L\* comprises a carbonyl linkage to said hydroxy residue.

8 A compound according to claim 7, wherein L\*' further comprises an  
25 intermediate linking moiety possessing an hydroxy group to which Rx is esterified.

9. A compound according to claim 8 wherein the intermediate linking moiety is an optionally branched, optionally monounsaturated alkylenoxy group having 1-3 chain carbons.

5 10 A compound according to claim 9 wherein L\*-Rx has the configuration:



wherein Ry and Ry' are independently H or C<sub>1</sub>-C<sub>3</sub> alkyl and PETT together with the arrowed oxygen represents the hydroxy residue of the compound of claim 1.

10

11 A compound according to claim 10, wherein Ry and Ry' are both hydrogen.

12 A compound according to claim 1, wherein Rx is derived from an  
15 aliphatic amino acid selected from alanine, leucine, isoleucine, tertiary leucine and valine.


13 A compound according to claim 12 wherein Rx is a derivative of an L-amino acid.

20

14 A compound according to claim 13 wherein Rx is a derivative of L-valine or L-isoleucine.

15. A compound according to claim 1, with the formula P2

25



wherein

R8 is halo;

R9 is C<sub>1</sub>-C<sub>3</sub> alkyl;

5 R10 is halo or cyano.

16 A compound according to claim 15, wherein R8 is fluoro.

17 A compound according to claim 15, wherein R9 is ethyl.


10

18 A compound according to claim 15, wherein R10 is cyano.

19 A compound according to claim 15, comprising at least 90% enantiomerically pure 1S 2S *cis* cyclopropyl configuration.

15

20 A compound according to claim 15, with the formula P3



20 wherein


Ry and Ry' are independently H or C<sub>1-3</sub> alkyl.

21 A compound according to claim 20, denoted:

(1S, 2S)-N-[*cis*-2-(6-fluoro, 2-(L-valyloxymethoxy), 3-propionylphenyl)-cyclopropyl]-N'-(5-cyanopyrid-2-yl)-urea or (1S, 2S)-N-[*cis*-2-(6-fluoro, 2-(L-isoleucyloxymethoxy), 3-propionylphenyl)-cyclopropyl]-N'-(5-bromopyrid-2-yl)-urea.

5

22 A compound according to claim 15 with the formula:



wherein

10 Alk<sup>b</sup> is an optionally branched, optionally unsaturated C<sub>1-6</sub> alkyl,

Alk is C<sub>1-6</sub> alkyl,

T is a bond or -O-

Rx' is the ester residue of a natural or unnatural amino acid, which may be the same or different to Rx;

15 and k is 0 or 1.

23 A compound according to claim 22, wherein Ry and Ry' are each H.

24 A compound according to claim 22, wherein R<sub>2</sub> is derived from L-valyl or  
20 L-isoleucyl.

25 A compound according to claim 22, wherein T is -O-.

26 A compound according to claim 22, wherein T is a bond.

25

27 A compound according to claim 22, wherein Alk<sup>b</sup> is 1,1 dimethylethylene, 2,2-dimethylethylene, propylene or butylene.

28 A compound according to claim 22 wherein k is 0.

29 A compound according to claim 22 denoted

(1S,2S)-N-[*cis*-2-(6-fluoro-2-(1,3-bis-L-valyloxy-2-

(oxycarbonylmethoxy)propyl)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,

(1S,2S)-N-[*cis*-2-(6-fluoro-2-(1,3-bis-L-isoleucyloxy-2-(oxycarbonylmethoxy)propyl)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,

(1S,2S)-N-[*cis*-2-(6-fluoro-2-(1,3-bis-L-valyloxy-2-

(oxycarbonylmethoxy)propyl)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,

(1S,2S)-N-[*cis*-2-(6-fluoro-2-(1,3-bis-L-isoleucyloxy-2-(oxycarbonylmethoxy)propyl)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,

(1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2-(L-valyloxy)-ethoxycarbonyloxymethyloxy)-3-propionylphenyl]]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,


(1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2-(L-isoleucyloxy)-ethoxycarbonyloxymethyloxy)-3-propionylphenyl]]cyclopropyl}-N'-[2-(5-cyanopyridyl)] urea,

(1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2-(L-valyloxy)-ethoxycarbonyloxymethyloxy)-3-propionylphenyl]]cyclopropyl}-N'-[2-(5-bromopyridyl)] urea,

(1S, 2S)-N-{*cis*-2-[6-fluoro-2-(2-(L-isoleucyloxy)-ethoxycarbonyloxymethyloxy)-3-propionylphenyl]]cyclopropyl}-N'-[2-(5-bromocyanopyridyl)] urea,

25

30 A compound according to claim 10 with the formula



wherein Alk<sup>b</sup> is optionally branched, optionally monounsaturated C<sub>1</sub>-C<sub>6</sub> alkyl.

31 A compound according to claim 30, wherein Alk<sup>b</sup> is methylene or , ethylene.

- 5 32 A compound according to claim 30 denoted  
(1S,2S)-N-[*cis*-2-(6-fluoro-2-(L-valyloxymethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,  
(1S,2S)-N-[*cis*-2-(6-fluoro-2-(L-isoleucyloxymethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,  
10 (1S,2S)-N-[*cis*-2-(6-fluoro-2-(L-valyloxymethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,  
(1S,2S)-N-[*cis*-2-(6-fluoro-2-(L-isoleucyloxymethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,  
(1S,2S)-N-[*cis*-2-(6-fluoro-2-(2-(L-valyloxy)ethoxycarbonyloxy)-3-  
15 propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,  
(1S,2S)-N-[*cis*-2-(6-fluoro-2-(2-(L-isoleucyloxy)ethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,  
(1S,2S)-N-[*cis*-2-(6-fluoro-2-(2-(L-valyloxy)ethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,  
20 (1S,2S)-N-[*cis*-2-(6-fluoro-2-(2-L-isoleucyloxy)ethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,  
(1S,2S)-N-[*cis*-2-(6-fluoro-2-(3-(L-valyloxy)propoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,  
(1S,2S)-N-[*cis*-2-(6-fluoro-2-(3-(L-isoleucyloxy)propoxycarbonyloxy)-3-  
25 propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,  
(1S,2S)-N-[*cis*-2-(6-fluoro-2-(3-(L-valyloxy)propoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,  
(1S,2S)-N-[*cis*-2-(6-fluoro-2-(3-(L-isoleucyloxy)propoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,  
30 (1S,2S)-N-[*cis*-2-(6-fluoro-2-(4-(L-valyloxy)butoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,  
(1S,2S)-N-[*cis*-2-(6-fluoro-2-(4-(L-isoleucyloxy)butoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea,

- (1S,2S)-N-[*cis*-2-(6-fluoro-2-(4-(L-valyloxy)butoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,  
(1S,2S)-N-[*cis*-2-(6-fluoro-2-(4-(L-isoleucyloxy)butoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-bromopyridyl)]urea,  
5 or a pharmaceutically acceptable salt thereof.

33. A compound according to claim 32 denoted (1S,2S)-N-[*cis*-2-(6-fluoro-2-(L-valyloxymethoxycarbonyloxy)-3-propionylphenyl)cyclopropyl]-N'-[2-(5-cyanopyridyl)]urea or a pharmaceutically acceptable salt thereof.

10

34 A pharmaceutical composition comprising a compound as claimed in claim 1 and a pharmaceutically acceptable carrier therefor.

35. A pharmaceutical composition according to claim 34, further comprising  
15 one to three additional antivirals selected from AZT, ddI, ddC, D4T, 3TC, DAPD, abacavir, adefovir, adefovir dipivoxil, bis-POC-PMPA, emtricitabine, tamociclovir (H2G), valtamociclovir stearate (MIV-606), hydroxyurea, Hoechst-Bayer HBY 097, foscarnet (PFA), efavirenz, trovirdine, nevirapine, delaviridine, emivirine, DMP-450, loviride, ritonavir, ABT 378, saquinavir,  
20 indinavir, amprenavir (Vertex VX 478) and nelfinavir.

36 A compound as claimed in claim 1, for use in therapy.

37 Use of a compound as claimed in claim 1 in the preparation of a  
25 medicament for the treatment or prophylaxis of HIV infection.

38 A method for the treatment or prophylaxis of HIV in a subject in need thereof comprising the administration of an effective amount of a compound as defined in claim 1 to the subject.

# INTERNATIONAL SEARCH REPORT

Int ional Application No  
PCT/SE 99/01403

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C07D213/85 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 99 41275 A (MEDIVIR AB) 19 August 1999 (1999-08-19) examples 12-19	1-38
X,P	WO 99 36406 A (SAHLBERG CHRISTER ; NOREEN ROLF (SE); ENGELHARDT PER (SE); HOEGBERG) 22 July 1999 (1999-07-22) cited in the application claims 6,22-27; examples	1,34-38
A	WO 95 06034 A (MEDIVIR AB ; LIND PETER THOMAS (SE); NOREEN ROLF (SE); MORIN JOHN M) 2 March 1995 (1995-03-02) cited in the application claims 1,15-18; examples	1,34-38
-/-		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "B" document member of the same patent family

Date of the actual completion of the international search  3 December 1999	Date of mailing of the international search report  11/01/2000
Name and mailing address of the ISA European Patent Office, P.B. 5618 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 eponi, Fax (+31-70) 340-3018	Authorized officer  Bosma, P



# INTERNATIONAL SEARCH REPORT

Int. Patent Application No.  
PCT/SE 99/01403

## C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BELL ET AL: "Phenethylthiazolethiourea (PETT) Compounds, a New Class of HIV-1 Reverse Transcriptase Inhibitors. 1. Synthesis and Basic Structure-Activity Relationship Studies of PETT Analogs" JOURNAL OF MEDICINAL CHEMISTRY, US, AMERICAN CHEMICAL SOCIETY. WASHINGTON, vol. 38, no. 25, page 4261-4274 XP002108200 ISSN: 0022-2623 cited in the application tables 1,2</p>	1, 34-38

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 99/01403

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 38  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claim 38  
is directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Int lional Application No

PCT/SE 99/01403

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9941275 A	19-08-1999	AU 8754898 A	08-03-1999
		WO 9909031 A	25-02-1999
		WO 9941268 A	19-08-1999
WO 9936406 A	22-07-1999	AU 2445099 A	02-08-1999
WO 9506034 A	02-03-1995	AT 173466 T	15-12-1998
		AU 687440 B	26-02-1998
		AU 7715394 A	21-03-1995
		CA 2168447 A	02-03-1995
		DE 69414708 D	24-12-1998
		DE 69414708 T	15-04-1999
		EP 0706514 A	17-04-1996
		ES 2123156 T	01-01-1999
		GR 3029370 T	28-05-1999
		JP 9502702 T	18-03-1997

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record.**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**